

The Lie algebra structure on the first Hochschild cohomology group of a monomial algebra

Claudia Strametz

Département de Mathématiques, Université de Montpellier 2,

F-34095 Montpellier cedex 5, France

E-mail: strametz@math.univ-montp2.fr

Résumé

Nous étudions la structure d'algèbre de Lie du premier groupe de la cohomologie de Hochschild d'une algèbre monomiale de dimension finie Λ , en termes combinatoires de son carquois, en quelconque caractéristique. Cela nous permet aussi d'examiner la composante de l'identité du groupe algébrique des automorphismes extérieurs de Λ en caractéristique zéro. Nous donnons des critères pour la (semi-)simplicité la résolubilité.

Abstract

We study the Lie algebra structure of the first Hochschild cohomology group of a finite dimensional monomial algebra Λ , in terms of the combinatorics of its quiver, in any characteristic. This allows us also to examine the identity component of the algebraic group of outer automorphisms of Λ in characteristic zero. Criteria for the (semi-)simplicity, the solvability, the reductivity, the commutativity and the nilpotency are given.

2000 Mathematics Subject Classification: 16E40 16W20

Keywords: cohomology, Hochschild, Lie, monomial

1 Introduction

The Hochschild cohomology $H^*(\Lambda, \Lambda)$ of any associative algebra Λ over a field k has the structure of a Gerstenhaber algebra (see [5]). In particular, the first Hochschild cohomology group $H^1(\Lambda, \Lambda)$ is a Lie algebra, a fact which can be verified directly. Note that in the finite dimensional case in characteristic 0 this Lie algebra can be regarded as the Lie algebra of the algebraic group of outer automorphisms $\text{Out}(\Lambda) = \text{Aut}(\Lambda)/\text{Inn}(\Lambda)$ of the algebra Λ . It has been treated by Guil-Asensio and Saorín in [8]. Huisgen-Zimmermann and Saorín proved in [11] that the identity component $\text{Out}(\Lambda)^\circ$ of the outer automorphism group of Λ is invariant under derived equivalence and Keller [13] showed that Hochschild cohomology is preserved under derived equivalence as a graded (super) Lie algebra. Consequently the Lie algebra $H^1(\Lambda, \Lambda)$ is invariant under derived equivalence.

The purpose of this paper is to study the Lie algebra structure of $H^1(\Lambda, \Lambda)$ in the case of finite dimensional monomial algebras without any restriction on the characteristic of the field k using only algebraic methods. The relationship between $H^1(\Lambda, \Lambda)$ and $\text{Out}(\Lambda)$ allows us to transfer the results obtained in this way to the identity component $\text{Out}(\Lambda)^\circ$ of the algebraic group $\text{Out}(\Lambda)$ in characteristic

0. Thus we give a different proof of Guil-Asensio and Saorín's criterion for the solvability of $\text{Out}(\Lambda)^\circ$ and we generalize some results they obtained using algebraic group theory and methods in algebraic geometry.

This paper is organized in the following way: in the first section we will use the minimal projective resolution of a monomial algebra Λ (as a Λ -bimodule) given by Bardzell in [1] to get a handy description of the Lie algebra $H^1(\Lambda, \Lambda)$ in terms of parallel paths. The purpose of the second section is to link this description to Guil-Asensio and Saorín's work on the algebraic group of outer automorphisms of monomial algebras (see [9]). In section three we carry out the study of the Lie algebra $H^1(\Lambda, \Lambda)$. In particular we give criteria for the (semi-)simplicity, the solvability, the reductivity, the commutativity and the nilpotency of this Lie algebra and consequently of the connected algebraic group $\text{Out}(\Lambda)^\circ$, in terms of the combinatorics of the quiver of Λ . Finally a Morita equivalence given by Gabriel in [3] in case $\text{char } k = p > 0$ between group algebras kG where the finite group G admits a normal cyclic Sylow p -subgroup and certain monomial algebras allows us to apply some of our results to the Lie algebra $H^1(kG, kG)$. Note that the Hochschild cohomology is Morita invariant as a Gerstenhaber algebra (see [6]).

We give some notation and terminology which we keep throughout the paper. Let Q denote a finite quiver (that is a finite oriented graph) and k an algebraically closed field. For all $n \in \mathbb{N}$ let Q_n be the set of oriented paths of length n of Q . Note that Q_0 is the set of vertices and that Q_1 is the set of arrows of Q . We denote by $s(\gamma)$ the source vertex of an (oriented) path γ of Q and by $t(\gamma)$ its terminus vertex. The path algebra kQ is the k -linear span of the set of paths of Q where the multiplication of $\beta \in Q_i$ and $\alpha \in Q_j$ is provided by the concatenation $\beta\alpha \in Q_{i+j}$ if $t(\alpha) = s(\beta)$ and 0 otherwise.

We denote by Λ a finite dimensional monomial k -algebra, that is a finite dimensional k -algebra which is isomorphic to a quotient of a path algebra kQ/I where the two-sided ideal I of kQ is generated by a set Z of paths of length ≥ 2 . We shall assume that Z is minimal, *i.e.* no proper subpath of a path in Z is again in Z . Let B be the set of paths of Q which do not contain any element of Z as a subpath. It is clear that the (classes modulo $I = \langle Z \rangle$) of elements of B form a basis of Λ . We shall denote by B_n the subset $Q_n \cap B$ of B formed by the paths of length n .

Let $E \simeq kQ_0$ be the separable subalgebra of Λ generated by the (classes modulo I of the) vertices of Q . We have a Wedderburn-Malcev decomposition $\Lambda = E \oplus r$ where r denotes the Jacobson radical of Λ .

This work will form part of a Ph.D. thesis under the supervision of Claude Cibils. I would like to thank him for his comments and his encouragement which I appreciated.

2 Projective resolutions and the Lie bracket

The Hochschild cohomology $H^*(\Lambda, \Lambda) = \text{Ext}_{\Lambda^e}^*(\Lambda, \Lambda)$ of a k -algebra Λ can be computed using different projective resolutions of Λ over its enveloping algebra $\Lambda^e = \Lambda \otimes_k \Lambda^{\text{op}}$. The standard resolution $\mathcal{P}_{\text{Hoch}}$ is

$$\cdots \rightarrow \Lambda \otimes_k^n \xrightarrow{\delta} \Lambda \otimes_k^{n-1} \rightarrow \cdots \rightarrow \Lambda \otimes_k \Lambda \xrightarrow{\varepsilon} \Lambda \rightarrow 0$$

where $\varepsilon(x_1 \otimes_k x_2) = x_1 x_2$ and

$$\delta(x_1 \otimes_k \cdots \otimes_k x_n) = \sum_{i=1}^{n-1} (-1)^{i+1} x_1 \otimes_k \cdots \otimes_k x_i x_{i+1} \otimes_k \cdots \otimes_k x_n$$

for $x_1, \dots, x_n \in \Lambda$. Applying the functor $\text{Hom}_{\Lambda^e}(-, \Lambda)$ to $\mathcal{P}_{\text{Hoch}}$ and identifying $\text{Hom}_{\Lambda^e}(\Lambda \otimes_k \Lambda^{\otimes_k^n} \otimes_k \Lambda, \Lambda)$ with $\text{Hom}_k(\Lambda^{\otimes_k^n}, \Lambda)$ for all $n \in \mathbb{N}$, yields the cochain

complex $\mathcal{C}_{\text{Hoch}}$ defined by Hochschild:

$$0 \rightarrow \Lambda \xrightarrow{d_0} \text{Hom}_k(\Lambda, \Lambda) \rightarrow \cdots \rightarrow \text{Hom}_k(\Lambda^{\otimes_k^n}, \Lambda) \xrightarrow{d_n} \text{Hom}_k(\Lambda^{\otimes_k^{n+1}}, \Lambda) \rightarrow \cdots$$

where $(d_0 a)(x) = ax - xa$ for all $a, x \in \Lambda$ and

$$\begin{aligned} (d_n f)(x_1 \otimes_k \cdots \otimes_k x_{n+1}) &= x_1 f(x_2 \otimes_k \cdots \otimes_k x_{n+1}) + \\ &\quad \sum_{i=1}^n (-1)^i f(x_1 \otimes_k \cdots \otimes_k x_i x_{i+1} \otimes_k \cdots \otimes_k x_{n+1}) \\ &\quad + (-1)^{n+1} f(x_1 \otimes_k \cdots \otimes_k x_n) x_{n+1} \end{aligned}$$

for all $f \in \text{Hom}_k(\Lambda^{\otimes_k^n}, \Lambda)$, $n \in \mathbb{N}$, and $x_1, \dots, x_{n+1} \in \Lambda$. In 1962 the structure of a Gerstenhaber algebra was introduced on the Hochschild cohomology $H^*(\Lambda, \Lambda)$ by Gerstenhaber in [5]. In particular the first cohomology group $H^1(\Lambda, \Lambda)$ which is the quotient of the derivations modulo the inner derivations of Λ is a Lie algebra whose bracket is induced by the Lie bracket on $\text{Hom}_k(\Lambda, \Lambda)$

$$[f, g] = f \circ g - g \circ f$$

where $f, g \in \text{Hom}_k(\Lambda, \Lambda)$. The 1-coboundaries, *i.e.* the derivations $\text{Der}_k(\Lambda) = \text{Ker } d_1 = \{f \in \text{Hom}_k(\Lambda, \Lambda) \mid f(ab) = af(b) + f(a)b \forall a, b \in \Lambda\}$ form a Lie subalgebra of $\text{Hom}_k(\Lambda, \Lambda)$ and the 1-cocycles, *i.e.* the inner derivations $\text{Ad}_k(\Lambda) = \text{Im } d_0 = \{f \in \text{Hom}_k(\Lambda, \Lambda) \mid \exists a \in \Lambda \text{ such that } f(x) = ax - xa \forall x \in \Lambda\}$ an ideal of this Lie algebra.

In order to compute the first Hochschild cohomology group $H^1(\Lambda, \Lambda)$ of the finite dimensional monomial algebra Λ , we shall use the minimal projective resolution of the Λ -bimodule Λ given by Bardzell in [1]. The part of this resolution $\mathcal{P}_{\min}$ in which we are interested is the following:

$$\cdots \rightarrow \Lambda \otimes_E kZ \otimes_E \Lambda \xrightarrow{\delta_1} \Lambda \otimes_E kQ_1 \otimes_E \Lambda \xrightarrow{\delta_0} \Lambda \otimes_E \Lambda \xrightarrow{\pi} \Lambda \rightarrow 0$$

where the Λ -bimodule morphisms are given by

$$\begin{aligned} \pi(\lambda \otimes_E \mu) &= \lambda \mu \\ \delta_0(\lambda \otimes_E a \otimes_E \mu) &= \lambda a \otimes_E \mu - \lambda \otimes_E a \mu \quad \text{and} \\ \delta_1(\lambda \otimes_E p \otimes_E \mu) &= \sum_{d=1}^n \lambda a_n \cdots a_{d+1} \otimes_E a_d \otimes_E a_{d-1} \cdots a_1 \mu \end{aligned}$$

for all $\lambda, \mu \in \Lambda$, $a, a_n, \dots, a_1 \in Q_1$ and $p = a_n \cdots a_1 \in Z$ (with the conventions $a_{n+1} = t(a_n)$ and $a_0 = s(a_1)$).

Remark 2.1 Our description of $\mathcal{P}_{\min}$ is equivalent to Bardzell's, because if X denotes a set of paths and if kX is the corresponding E -bimodule, then the map $\oplus_{\gamma \in X} \Lambda t(\gamma) \otimes_k s(\gamma) \Lambda \rightarrow \Lambda \otimes_E kX \otimes_E \Lambda$ which is given by $\lambda t(\gamma) \otimes_k s(\gamma) \mu \mapsto \lambda \otimes_E \gamma \otimes_E \mu$, where $\gamma \in X$ and $\lambda, \mu \in \Lambda$, is clearly a Λ -bimodule isomorphism. Note that the Λ -bimodules $\Lambda \otimes_E \Lambda$ and $\Lambda \otimes_E E \otimes_E \Lambda \simeq \Lambda \otimes_E kQ_0 \otimes_E \Lambda$ are isomorphic.

Lemma 2.2 Let M be an E -bimodule and T a Λ -bimodule. Then the vector space $\text{Hom}_{\Lambda^e}(\Lambda \otimes_E M \otimes_E \Lambda, T)$ is isomorphic to $\text{Hom}_{E^e}(M, T)$.

Proof: The linear morphism $\text{Hom}_{\Lambda^e}(\Lambda \otimes_E M \otimes_E \Lambda, T) \rightarrow \text{Hom}_{E^e}(M, T)$ which sends a Λ -bimodule morphism $f : \Lambda \otimes_E M \otimes_E \Lambda \rightarrow T$ to the E -bimodule morphism $M \rightarrow T$ given by $m \mapsto f(1_\Lambda \otimes_E m \otimes_E 1_\Lambda)$ and the linear morphism $\text{Hom}_{E^e}(M, T) \rightarrow \text{Hom}_{\Lambda^e}(\Lambda \otimes_E M \otimes_E \Lambda, T)$ which associates to $g : M \rightarrow T$ the element $\Lambda \otimes_E M \otimes_E \Lambda \rightarrow T$, $\lambda \otimes_E m \otimes_E \mu \mapsto \lambda g(m) \mu$, are inverse to each other.

q.e.d. 2.2

Applying the functor $\text{Hom}_{\Lambda^e}(-, \Lambda)$ to $\mathcal{P}_{\min}$ and using the preceding Lemma yields the cochain complex $\mathcal{C}_{\min}$

$$0 \longrightarrow \text{Hom}_{E^e}(kQ_0, \Lambda) \xrightarrow{\delta_0^*} \text{Hom}_{E^e}(kQ_1, \Lambda) \xrightarrow{\delta_1^*} \text{Hom}_{E^e}(kZ, \Lambda) \longrightarrow \dots$$

where the coboundaries δ_0^* and δ_1^* are given by

$$\begin{aligned} (\delta_0^* f)(a) &= af(s(a)) - f(t(a))a & \text{and} \\ (\delta_1^* g)(p) &= \sum_{d=1}^n a_n \dots a_{d+1} g(a_d) a_{d-1} \dots a_1 \end{aligned}$$

where $f \in \text{Hom}_{E^e}(kQ_0, \Lambda)$, $a, a_n, \dots, a_1 \in Q_1$, $g \in \text{Hom}_{E^e}(kQ_1, \Lambda)$ and $p = a_n \dots a_1 \in Z$.

As it is useful to interpret $H^1(\Lambda, \Lambda)$ in terms of parallel paths we introduce now the following notion: two paths ε, γ of Q are called parallel if $s(\varepsilon) = s(\gamma)$ and $t(\varepsilon) = t(\gamma)$. If X and Y are sets of paths of Q , the set $X//Y$ of parallel paths is formed by the couples $(\varepsilon, \gamma) \in X \times Y$ such that ε and γ are parallel paths. For instance, $Q_0//Q_n$ is the set of oriented cycles of Q of length n .

Lemma 2.3 *Let X and Y be sets of paths of Q and let kX and kY be the corresponding E -bimodules. Then the vector spaces $k(X//Y)$ and $\text{Hom}_{E^e}(kX, kY)$ are isomorphic.*

Proof: Define a linear morphism $k(X//Y) \rightarrow \text{Hom}_{E^e}(kX, kY)$ by sending $(\varepsilon, \gamma) \in X//Y$ to the elementary map which associates γ to ε and 0 to any other path of X . If $f : kX \rightarrow kY$ is an E -bimodule morphism, we have for every path $\varepsilon \in X$ that $f(\varepsilon) = f(t(\varepsilon)\varepsilon s(\varepsilon)) = t(\varepsilon)f(\varepsilon)s(\varepsilon)$ and thus $f(\varepsilon) = \sum_{(\varepsilon, \gamma) \in X//Y} \lambda_{\varepsilon, \gamma} \gamma$, where $\lambda_{\varepsilon, \gamma} \in k$. This allows us to define a linear morphism $\text{Hom}_{E^e}(kX, kY) \rightarrow k(X//Y)$ which assigns to $f : kX \rightarrow kY$ the element $\sum_{(\varepsilon, \gamma) \in X//Y} \lambda_{\varepsilon, \gamma} (\varepsilon, \gamma)$. Obviously, the two morphisms are inverse to each other.

q.e.d. 2.3

Using the following notations we will rewrite the coboundaries:

Notations 2.4

- (i) Let ε be a path in Q and $(a, \gamma) \in Q_1//B$. We denote by $\varepsilon^{(a, \gamma)}$ the sum of all nonzero paths (i.e. paths in B) obtained by replacing one appearance of the arrow a in ε by the path γ . If the path ε does not contain the arrow a or if every replacement of a in ε by γ is not a path in B , we set $\varepsilon^{(a, \gamma)} = 0$. Suppose that $\varepsilon^{(a, \gamma)} = \sum_{i=1}^n \varepsilon_i$, where $\varepsilon_i \in B$ and let η be a path of B parallel to ε . By abuse of language we denote by $(\eta, \varepsilon^{(a, \gamma)})$ the sum $\sum_{i=1}^n (\eta, \varepsilon_i)$ in $k(Z//B)$ (with the convention that $(\eta, \varepsilon^{(a, \gamma)}) = 0$ if $\varepsilon^{(a, \gamma)} = 0$).
- (ii) The function $\chi_B : \prod_{n \in \mathbb{N}} Q_n \rightarrow \{0, 1\}$ denotes the indicator function which associates 1 to each path $\gamma \in B$ and 0 to $\gamma \notin B$.
- (iii) If X is a set of paths of Q and e a vertex of Q , the set Xe is formed by the paths of X with source vertex e . In the same way eX denotes the set of all paths of X with terminus vertex e .

Remark 2.5

If $(a, \gamma) \in Q_1//(B - Q_0)$, then all the nonzero summands of $\varepsilon^{(a, \gamma)}$ are distinct.

If we carry out the identification suggested in the preceding Lemma, we obtain:

Proposition 2.6

The beginning of the cochain complex $\mathcal{C}_{\min}$ can be characterized in the following way

$$0 \rightarrow k(Q_0//B) \xrightarrow{\psi_0} k(Q_1//B) \xrightarrow{\psi_1} k(Z//B) \xrightarrow{\psi_2} \dots$$

where the maps are given by

$$\begin{aligned} \psi_0 : \quad k(Q_0//B) &\longrightarrow k(Q_1//B) \\ (e, \gamma) &\longmapsto \sum_{a \in Q_1 e} \chi_B(a\gamma)(a, a\gamma) - \sum_{a \in e Q_1} \chi_B(\gamma a)(a, \gamma a) \\ \\ \psi_1 : \quad k(Q_1//B) &\longrightarrow k(Z//B) \\ (a, \gamma) &\longmapsto \sum_{p \in Z} (p, p^{(a, \gamma)}) \end{aligned}$$

In particular, we have $H^1(\Lambda, \Lambda) \simeq \text{Ker } \psi_1 / \text{Im } \psi_0$.

The verifications are straightforward.

Theorem 2.7 *The bracket*

$$[(a, \gamma), (b, \varepsilon)] = (b, \varepsilon^{(a, \gamma)}) - (a, \gamma^{(b, \varepsilon)})$$

for all $(a, \gamma), (b, \varepsilon) \in Q_1//B$ induces a Lie algebra structure on $\text{Ker } \psi_1 / \text{Im } \psi_0$ such that $H^1(\Lambda, \Lambda)$ and $\text{Ker } \psi_1 / \text{Im } \psi_0$ are isomorphic Lie algebras.

Proof: As $\mathcal{P}_{\text{Hoch}}$ and $\mathcal{P}_{\min}$ are projective resolutions of the Λ -bimodule Λ , there exist, thanks to the Comparison Theorem, chain maps $\omega : \mathcal{P}_{\text{Hoch}} \rightarrow \mathcal{P}_{\min}$ and $\varsigma : \mathcal{P}_{\min} \rightarrow \mathcal{P}_{\text{Hoch}}$. Let us choose these Λ -bimodule morphisms such that

$$\begin{aligned} \varsigma_0 : \quad & \Lambda \otimes_E \Lambda \longrightarrow \Lambda \otimes_k \Lambda \\ & \lambda \otimes_E \mu \longmapsto \lambda \otimes_k \mu \\ \varsigma_1 : \quad & \Lambda \otimes_E kQ_1 \otimes_E \Lambda \longrightarrow \Lambda \otimes_k \Lambda \otimes_k \Lambda \\ & \lambda \otimes_E a \otimes_E \mu \longmapsto \lambda \otimes_k a \otimes_k \mu \\ \omega_0 : \quad & \Lambda \otimes_k \Lambda \longrightarrow \Lambda \otimes_E \Lambda \\ & \lambda \otimes_k \mu \longmapsto \lambda \otimes_E \mu \\ \omega_1 : \quad & \Lambda \otimes_k \Lambda \otimes_k \Lambda \longrightarrow \Lambda \otimes_E kQ_1 \otimes_E \Lambda \\ & \lambda \otimes_k a_n \dots a_1 \otimes_k \mu \longmapsto \sum_{d=1}^n \lambda a_n \dots a_{d+1} \otimes_E a_d \otimes_E a_{d-1} \dots a_1 \mu \end{aligned}$$

where $\lambda, \mu \in \Lambda$, $a, a_n, \dots, a_1 \in Q_1$ and $a_n \dots a_1 \in B$. As a chain map between projective resolutions is unique up to chain homotopy equivalence, ω and ς are unique up to chain homotopy equivalence and such that $\omega \circ \varsigma$ is homotopic to $\text{id}_{\mathcal{P}_{\min}}$ and $\varsigma \circ \omega$ is homotopic to $\text{id}_{\mathcal{P}_{\text{Hoch}}}$. Therefore, the cochain maps $\text{Hom}_{\Lambda^e}(\omega, \Lambda)$ and $\text{Hom}_{\Lambda^e}(\varsigma, \Lambda)$ are such that $\text{Hom}_{\Lambda^e}(\varsigma, \Lambda) \circ \text{Hom}_{\Lambda^e}(\omega, \Lambda)$ is homotopic to $\text{id}_{\mathcal{C}_{\min}}$ and $\text{Hom}_{\Lambda^e}(\omega, \Lambda) \circ \text{Hom}_{\Lambda^e}(\varsigma, \Lambda)$ is homotopic to $\text{id}_{\text{Hom}_{\Lambda^e}(\mathcal{P}_{\text{Hoch}}, \Lambda)}$. Hence these maps induce inverse linear isomorphisms at the cohomology level. Taking into account the identifications above we obtain that

$$\begin{aligned} \overline{\omega}_1 : \quad & k(Q_1//B) \longrightarrow \text{Hom}_k(\Lambda, \Lambda) \\ & (a, \gamma) \longmapsto \begin{array}{ccc} \Lambda & \longrightarrow & \Lambda \\ \varepsilon & \longmapsto & \varepsilon^{(a, \gamma)} \end{array} \quad \text{and} \\ \\ \overline{\varsigma}_1 : \quad & \text{Hom}_k(\Lambda, \Lambda) \longrightarrow k(Q_1//B) \\ f : \Lambda &\longrightarrow \Lambda \longmapsto \sum_{a \in Q_1} \sum_{(a, \gamma) \in Q_1//B} \lambda_{a, \gamma}(a, \gamma) \\ \varepsilon &\longmapsto \sum_{\gamma \in B} \lambda_{\varepsilon, \gamma} \gamma \end{aligned}$$

induce inverse linear isomorphisms between $H^1(\Lambda, \Lambda) = \text{Ker } d_1 / \text{Im } d_0$ and $H^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$. This allows us to transfer the Lie algebra structure of $\text{Ker } d_1 / \text{Im } d_0$ to $\text{Ker } \psi_1 / \text{Im } \psi_0$. Define on $k(Q_1//B)$ the bracket

$$[(a, \gamma), (b, \varepsilon)] := \overline{\varsigma}_1([\overline{\omega}_1(a, \gamma), \overline{\omega}_1(b, \varepsilon)]) = (b, \varepsilon^{(a, \gamma)}) - (a, \gamma^{(b, \varepsilon)})$$

for all $(a, \gamma), (b, \varepsilon) \in Q_1//B$. It is easy to check that this is a Lie bracket. As $\overline{\omega} : \mathcal{C}_{\text{Hoch}} \rightarrow \mathcal{C}_{\text{min}}$ and $\overline{\tau} : \mathcal{C}_{\text{min}} \rightarrow \mathcal{C}_{\text{Hoch}}$ are maps of complexes, we conclude from the fact that $\text{Ker } d_1 = \text{Der}_k(\Lambda)$ is a Lie subalgebra of $\text{Hom}_k(\Lambda, \Lambda)$, that $\text{Ker } \psi_1$ is a Lie subalgebra of $k(Q_1//B)$. In the same way we deduce from the fact that $\text{Im } \psi_0 = \text{Ad}_k(\Lambda)$ is a Lie ideal of $\text{Ker } d_1$, that $\text{Im } \psi_0$ is a Lie ideal of $\text{Ker } \psi_1$. By construction the quotient Lie algebra $H^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$ is isomorphic to the Lie algebra $H^1(\Lambda, \Lambda) = \text{Ker } d_1 / \text{Im } d_0$.

q.e.d. 2.7

It is remarkable that Bardzell's minimal projective resolution and the resolution considered by Gerstenhaber and Schack in [7] using E -relative Hochschild cohomology yield the same 1-coboundaries and 1-cocycles as the following Proposition and Corollary show:

Proposition 2.8 *The Lie algebras $\text{Ker } \psi_1$ (with the bracket described in the preceding Theorem) and $\text{Der}_{E^e}(\Lambda)$ (with the canonical bracket) are isomorphic.*

Proof: Let $\sigma : \Lambda \rightarrow \Lambda$ be an element of $\text{Der}_{E^e}(\Lambda) = \text{Der}_k(\Lambda) \cap \text{Hom}_{E^e}(\Lambda, \Lambda)$. Since σ is an E -bimodule morphism, the image of an arrow a is a linear combination of paths of B parallel to a . Suppose $\sigma(a) = \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} \gamma$ for $a \in Q_1$ with $\lambda_{(a, \gamma)} \in k$. The fact that σ is a well-defined derivation implies that we have for every path $p = p_l \dots p_2 p_1$ in Z of length l

$$0 = \sigma(p) = \sigma(p_l \dots p_1) = \sum_{i=1}^l p_l \dots p_{i+1} \sigma(p_i) p_{i-1} \dots p_1 = \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} p^{(a, \gamma)}$$

Thus we have in $k(Z//B)$ the equality $0 = \sum_{p \in Z} \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} (p, p^{(a, \gamma)})$ which is the same as to say that $\sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} (a, \gamma)$ is an element in $\text{Ker } \psi_1$. This enables us to define the linear function

$$\xi : \text{Der}_{E^e}(\Lambda) \rightarrow \text{Ker } \psi_1, \quad \sigma \mapsto \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} (a, \gamma)$$

On the other hand we can associate to every $x = \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} (a, \gamma)$ in $\text{Ker } \psi_1$ an E -bimodule morphism $\sigma_x : \Lambda \rightarrow \Lambda$ by setting $\sigma_x(e) := 0$ for every vertex $e \in Q_0$ and $\sigma_x(c) := \sum_{(c, \gamma) \in Q_1//B} \lambda_{(c, \gamma)} \gamma$ for every arrow $c \in Q_1$. To make σ_x a derivation we extend σ_x to paths $\varepsilon = b_m \dots b_1$ of length $m \geq 2$ by the formula

$$\sigma_x(\varepsilon) = \sigma_x(b_m \dots b_1) = \sum_{j=1}^m b_m \dots b_{j+1} \sigma_x(b_j) b_{j-1} \dots b_1 = \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} \varepsilon^{(a, \gamma)}$$

Thanks to $0 = \psi_1(x) = \sum_{p \in Z} \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} (p, p^{(a, \gamma)})$ we have for every $p \in Z$ the relation $0 = \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} p^{(a, \gamma)} = \sigma_x(p)$ which shows that σ_x is well-defined. This allows us to define the linear function

$$\zeta : \text{Ker } \psi_1 \rightarrow \text{Der}_{E^e}(\Lambda), \quad x \mapsto \sigma_x$$

It is clear that ξ and ζ are inverse to each other which proves that $\text{Der}_{E^e}(\Lambda)$ and $\text{Ker } \psi_1$ are isomorphic k -vector spaces. In order to check that ζ is a Lie algebra morphism let us fix $x = \sum_{(a, \gamma) \in Q_1//B} \lambda_{(a, \gamma)} (a, \gamma)$ and $y = \sum_{(b, \varepsilon) \in Q_1//B} \mu_{(b, \varepsilon)} (b, \varepsilon)$

in $\text{Ker } \psi_1$. For every arrow $c \in Q_1$ we have

$$\begin{aligned}
[\sigma_x, \sigma_y](c) &= (\sigma_x \circ \sigma_y - \sigma_y \circ \sigma_x)(c) \\
&= \sum_{(c, \varepsilon) \in Q_1 // B} \mu_{(c, \varepsilon)} \sigma_x(\varepsilon) - \sum_{(c, \gamma) \in Q_1 // B} \lambda_{(c, \gamma)} \sigma_y(\gamma) \\
&= \sum_{(c, \varepsilon), (a, \gamma) \in Q_1 // B} \mu_{(c, \varepsilon)} \lambda_{(a, \gamma)} \varepsilon^{(a, \gamma)} - \sum_{(c, \gamma), (b, \varepsilon) \in Q_1 // B} \lambda_{(c, \gamma)} \mu_{(b, \varepsilon)} \gamma^{(b, \varepsilon)} \\
&= \sigma_z(c)
\end{aligned}$$

where $z = \sum_{(a, \gamma), (b, \varepsilon) \in Q_1 // B} \lambda_{(a, \gamma)} \mu_{(b, \varepsilon)} ((b, \varepsilon^{(a, \gamma)}) - (a, \gamma^{(b, \varepsilon)}))$. Theorem 2.7 shows that z is equal to the element $[x, y]$ of $\text{Ker } \psi_1$. Therefore ζ is a Lie algebra morphism and $\text{Der}_{E^e}(\Lambda)$ and $\text{Ker } \psi_1$ are isomorphic Lie algebras.

q.e.d. 2.8

Corollary 2.9 *The Lie ideal $\text{Im } \psi_0$ of $\text{Ker } \psi_1$ and the Lie ideal $\text{Ad}_{E^e}(\Lambda)$ of $\text{Der}_{E^e}(\Lambda)$ are isomorphic.*

Proof: Let $y = \sum_{\gamma \in B} \lambda_\gamma \gamma \in \Lambda$ be such that $\text{ad}(y)$ is an E -bimodule morphism. Then we have for every vertex $e \in Q_0$

$$0 = eye - eye = e \text{ad}(y)(e) = \text{ad}(y)(e) = ye - ey = \sum_{\gamma \in B} \lambda_\gamma (\delta_{s(\gamma), e} - \delta_{t(\gamma), e}) \gamma$$

This shows that every path $\gamma \in B$ satisfying $\lambda_\gamma \neq 0$ is a cycle and so $y = \sum_{(e, \gamma) \in Q_0 // B} \lambda_\gamma \gamma$. For every arrow $a \in Q_1$ we have

$$\text{ad}(y)(a) = ya - ay = \sum_{(e, \gamma) \in Q_0 // B} \lambda_\gamma \gamma a - \sum_{(e, \gamma) \in Q_0 // B} \lambda_\gamma a \gamma = \sum_{(e, \gamma) \in Q_0 // B} \lambda_\gamma (\gamma a - a \gamma)$$

We deduce from this that the image of $\text{ad}(y) \in \text{Ad}_{E^e}(\Lambda)$ by ξ is

$$\xi(\text{ad}(y)) = \sum_{(e, \gamma) \in Q_0 // B} \lambda_\gamma \left(\sum_{\substack{a \in e Q_1 \\ \gamma a \in B}} (a, \gamma a) - \sum_{\substack{a \in Q_1 e \\ a \gamma \in B}} (a, a \gamma) \right) = - \sum_{(e, \gamma) \in Q_0 // B} \lambda_\gamma \psi_0(e, \gamma)$$

and thus we obtain that the image of the Lie algebra $\text{Ad}_{E^e}(\Lambda)$ by the isomorphism of Lie algebras ξ is $\text{Im } \psi_0$.

q.e.d. 2.9

Remark 2.10 *In general, the minimal projective resolution and the resolution considered by Gerstenhaber and Schack do not yield the same n -coboundaries and n -cocycles: for example in the case of hereditary algebras, i.e. $Z = \emptyset$, we have for $n = 2$ that $\text{Ker } \psi_2 = 0$, because $k(Z // B) = 0$, but the multiplication $m \in \text{Hom}_{E^e}(\Lambda \otimes_E \Lambda, \Lambda)$ is a nontrivial 2-coboundary as well as a 2-cocycle of the complex yielded by Gerstenhaber and Schack's resolution.*

3 The algebraic group of outer automorphisms of a monomial algebra

In this section, we assume that the characteristic of the field k is 0. Our aim is to relate the description of the Lie algebra $\text{H}^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$ obtained in

the preceding section to the algebraic groups which appear in Guil-Asensio and Saorín's study of the outer automorphisms. Denote by $\text{Aut}(A)$ the algebraic group of all k -algebra automorphisms of a finite dimensional k -algebra A . The group of inner automorphisms $\text{Inn}(A)$ of A is the image of the morphism of algebraic groups $\varphi : A^* \rightarrow \text{Aut}(A)$ given by $a \mapsto \iota_a = \text{conjugation by } a$. Thus $\text{Inn}(A)$ is a closed normal and connected subgroup of $\text{Aut}(A)$. The algebraic group of outer automorphisms $\text{Out}(A)$ is defined as the quotient $\text{Aut}(A)/\text{Inn}(A)$. Its identity component $\text{Out}(A)^\circ$ is $\text{Aut}(A)^\circ/\text{Inn}(A)$. Note that if A is a basic algebra, for instance the monomial algebra Λ , then the group $\text{Out}(A)$ is isomorphic to the Picard group $\text{Pic}_k(A)$, that is the group of (isomorphism types of) Morita autoequivalences of the category of left A -modules (see [15], p. 1860).

Proposition 3.1 *Let k be a field of characteristic 0 and A a finite dimensional k -algebra. The derivations $\text{Der}_k(A)$ form the Lie algebra of the algebraic group $\text{Aut}(A)$ and the inner derivations $\text{Ad}_k(A)$ form the Lie algebra of the algebraic group $\text{Inn}(A)$. The Lie algebra $H^1(A, A)$ can be regarded as the Lie algebra of the algebraic group $\text{Out}(A)$ or as the Lie algebra of its identity component $\text{Out}(A)^\circ$.*

Proof: In [12] 13.2 it is shown that $\text{Der}_k(A)$ is the Lie algebra of $\text{Aut}(A)$. The differential $d\varphi : A \rightarrow \text{Der}_k(A)$ of the above-mentioned morphism φ is given by $a \mapsto \text{ad}(a) = \text{inner derivation of } a$. Since the field k has characteristic 0, the Lie algebra of the image of a morphism of algebraic groups is the image of the differential of this morphism. The construction of the quotient $\text{Aut}(A)/\text{Inn}(A)$ implies that the Lie algebra of the algebraic group $\text{Aut}(A)/\text{Inn}(A) = \text{Out}(A)$ is $\mathcal{L}(\text{Aut}(A))/\mathcal{L}(\text{Inn}(A)) = \text{Der}_k(A)/\text{Ad}_k(A) = H^1(A, A)$ (see [12] 11.5 and 12).

q.e.d. 3.1

The automorphism group of any finite dimensional algebra was studied by Pollack in [15]. Guil-Asensio and Saorín worked on the outer automorphisms of any finite dimensional algebra (see [8]). The case of the finite dimensional monomial algebras was treated by them in their paper [9]. We are going to follow their notations. Define the algebraic group

$$H_\Lambda := \{\sigma \in \text{Aut}(\Lambda) \mid \sigma(e) = e \ \forall e \in Q_0\}$$

and denote by H_Λ° its identity component. According to [15] and to Proposition 1.1 of [9] we have

$$\text{Out}(\Lambda)^\circ \simeq H_\Lambda^\circ/H_\Lambda \cap \text{Inn}(\Lambda)$$

Proposition 3.2 *Let k be a field of characteristic 0 and Λ a finite dimensional monomial algebra. The Lie algebra of the algebraic group H_Λ is the Lie algebra $\text{Der}_{E^e}(\Lambda) \simeq \text{Ker } \psi_1$ and $\text{Ad}_{E^e}(\Lambda) \simeq \text{Im } \psi_0$ is the Lie algebra of $H_\Lambda \cap \text{Inn}(\Lambda)$.*

Proof: For every vertex $e \in Q_0$ we write $G_e := \{\sigma \in \mathbf{GL}(\Lambda) \mid \sigma(e) = e\}$ and $\mathcal{G}_e := \{\sigma \in \mathfrak{gl}(\Lambda) \mid \sigma(e) = 0\}$. Thus we get the equality $H_\Lambda = \text{Aut}(\Lambda) \cap \bigcap_{e \in Q_0} G_e$. Since we have assumed $\text{char } k = 0$, paragraph 13.2 of [12] shows that

$$\begin{aligned} \mathcal{L}(H_\Lambda) &= \mathcal{L}(\text{Aut}(\Lambda)) \cap \bigcap_{e \in Q_0} \mathcal{L}(G_e) = \text{Der}_k(\Lambda) \cap \bigcap_{e \in Q_0} \mathcal{G}_e \\ &= \{\sigma \in \text{Der}_k(\Lambda) \mid \sigma(e) = 0 \ \forall e \in Q_0\} = \text{Der}_{E^e}(\Lambda) \quad \text{and} \end{aligned}$$

$$\mathcal{L}(H_\Lambda \cap \text{Inn}(\Lambda)) = \mathcal{L}(H_\Lambda) \cap \mathcal{L}(\text{Inn}(\Lambda)) = \text{Der}_{E^e}(\Lambda) \cap \text{Ad}_k(\Lambda) = \text{Ad}_{E^e}(\Lambda)$$

q.e.d. 3.2

The closed unipotent and connected subgroup

$$\hat{E} := \varphi(E^*) = \{\iota_a \in \text{Inn}(\Lambda) \mid a \in E^*\}$$

of H_Λ is isomorphic as a group to the group of acyclic characters $\text{Ch}(Q, k)$ appearing in Guil-Asensio and Saorín's paper (see definition 7 in [8]). The group

$$\text{Inn}^*(\Lambda) := \varphi(1 + r) = \{\iota_a \in \text{Inn}(\Lambda) \mid \exists x \in r \text{ such that } a = 1 + x\}$$

is a closed unipotent and connected subgroup of $\text{Inn}(\Lambda)$. We have $H_\Lambda \cap \text{Inn}(\Lambda) = (H_\Lambda \cap \text{Inn}^*(\Lambda)) \times \hat{E}$. Define the algebraic groups

$$U_\Lambda := \{\sigma \in \text{Aut}(\Lambda) \mid \sigma(a) \equiv a \pmod{r^2} \forall a \in Q_1\} \quad \text{and}$$

$$V_\Lambda := \{\sigma \in H_\Lambda \mid \sigma(a) = \sum_{(a,b) \in Q_1/Q_1} \lambda_{(a,b)} b \quad \forall a \in Q_1\}$$

According to Proposition 1.1 of [9] we have $H_\Lambda^\circ = (H_\Lambda \cap U_\Lambda) \rtimes V_\Lambda^\circ$ and therefore $\text{Out}(\Lambda)^\circ$ can be described as follows:

Proposition 3.3 *Let k be a field of characteristic 0 and Λ a finite dimensional monomial k -algebra. Then the identity component $\text{Out}(\Lambda)^\circ$ of the algebraic group of outer automorphisms is the semidirect product*

$$\text{Out}(\Lambda)^\circ = \frac{H_\Lambda \cap U_\Lambda}{H_\Lambda \cap \text{Inn}^*(\Lambda)} \rtimes \frac{V_\Lambda^\circ}{\hat{E}}$$

Since there is an inclusion preserving 1 – 1 correspondence between the collection of closed connected subgroups of $\text{Out}(\Lambda)^\circ$ (resp. H_Λ°) and the collection of their Lie algebras, regarded as subalgebras of $H^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$ (resp. $\text{Ker } \psi_1$) we are interested in identifying the subalgebras of $H^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$ (resp. $\text{Ker } \psi_1$) corresponding to the algebraic groups $\frac{H_\Lambda \cap U_\Lambda}{H_\Lambda \cap \text{Inn}^*(\Lambda)}$ and $\frac{V_\Lambda^\circ}{\hat{E}}$ (resp. $H_\Lambda \cap U_\Lambda$, $H_\Lambda \cap \text{Inn}^*(\Lambda)$, V_Λ° and $\hat{E}$). We have the following dictionary:

Proposition 3.4 *Let k be a field of characteristic 0 and Λ a finite dimensional monomial k -algebra.*

- (i) *The Lie algebra of the closed connected normal subgroup $H_\Lambda \cap U_\Lambda$ of H_Λ° is the Lie ideal $k(Q_1/B - (Q_0 \cup Q_1)) \cap \text{Ker } \psi_1$ of the Lie algebra $\text{Ker } \psi_1$.*
- (ii) *The Lie algebra of the closed connected subgroup V_Λ° of H_Λ° is the Lie subalgebra $k(Q_1/Q_1) \cap \text{Ker } \psi_1$ of the Lie algebra $\text{Ker } \psi_1$.*
- (iii) *The Lie algebra of the closed connected normal subgroup $H_\Lambda \cap \text{Inn}^*(\Lambda)$ of $H_\Lambda \cap \text{Inn}(\Lambda)$ is the Lie ideal of $\text{Im } \psi_0$ generated by the elements $\sum_{a \in Q_1} e(a, a\gamma) - \sum_{a \in eQ_1} (a, \gamma a)$ where $(e, \gamma) \in Q_0/(B - Q_0)$.*
- (iv) *The Lie algebra of the closed connected subgroup $\hat{E}$ of $H_\Lambda \cap \text{Inn}(\Lambda)$ is the Lie subalgebra of $\text{Im } \psi_0$ generated by the elements $\sum_{a \in Q_1} e(a, a) - \sum_{a \in eQ_1} (a, a)$ where $e \in Q_0$.*

Proof: The derivation of the morphism of algebraic groups

$$\begin{aligned} \varepsilon_\Lambda : \quad \text{Aut}(\Lambda) &\longrightarrow \mathbf{GL}(r/r^2) \simeq \mathbf{GL}(kQ_1) \\ \sigma &\longmapsto r/r^2 \rightarrow r/r^2, \quad x \pmod{r^2} \mapsto \sigma(x) \pmod{r^2} \end{aligned}$$

is given by

$$\begin{aligned} d\varepsilon_\Lambda : \quad \text{Der}_k(\Lambda) &\longrightarrow \mathfrak{gl}(r/r^2) \simeq \mathfrak{gl}(kQ_1) \\ \sigma &\longmapsto r/r^2 \rightarrow r/r^2, \quad x \pmod{r^2} \mapsto \sigma(x) \pmod{r^2} \end{aligned}$$

Recall that we have shown in Proposition 1.8 that

$$\begin{array}{lll} \xi : & \text{Der}_{E^e}(\Lambda) & \longrightarrow \text{Ker } \psi_1 \\ \sigma : & \Lambda \rightarrow \Lambda & \longmapsto \sum_{(a,\gamma) \in Q_1//B} \lambda_{(a,\gamma)}(a, \gamma) \\ a \in Q_1 & \mapsto \sum_{(a,\gamma) \in Q_1//B} \lambda_{(a,\gamma)}(a, \gamma) \end{array}$$

is a Lie algebra isomorphism.

(i) : Since U_Λ is the kernel of ε_Λ we obtain, thanks to the assumption $\text{char } k = 0$,

$$\mathcal{L}(H_\Lambda \cap U_\Lambda) = \mathcal{L}(H_\Lambda) \cap \text{Ker } d\varepsilon_\Lambda = \{\sigma \in \text{Der}_{E^e}(\Lambda) \mid \sigma(a) \in r^2 \ \forall a \in Q_1\}$$

The fact $\xi(\mathcal{L}(H_\Lambda \cap U_\Lambda)) = k(Q_1//B - (Q_0 \cup Q_1)) \cap \text{Ker } \psi_1$ finishes the proof.

(ii) : From $V_\Lambda \simeq \text{Im } \varepsilon_\Lambda|_{H_\Lambda}$ (see Lemma 22 in [8]) and $\text{char } k = 0$ we deduce

$$\begin{aligned} \mathcal{L}(V_\Lambda) &= \mathcal{L}(V_\Lambda^\circ) = \text{Im } d\varepsilon_\Lambda|_{\mathcal{L}(H_\Lambda)} \\ &= \{\sigma \in \text{Der}_{E^e}(\Lambda) \mid \sigma(a) = \sum_{(a,b) \in Q_1//Q_1} \lambda_{(a,b)}(a, b) \ \forall a \in Q_1\} \end{aligned}$$

The equality $\xi(\mathcal{L}(V_\Lambda^\circ)) = k(Q_1//Q_1) \cap \text{Ker } \psi_1$ shows that we are done.

(iii) : We have

$$\mathcal{L}(H_\Lambda \cap \text{Inn}^*(\Lambda)) = \{\text{ad}(1+x) \in \text{Ad}_{E^e}(\Lambda) \mid x \in r\} = \{\text{ad}(x) \in \text{Ad}_{E^e}(\Lambda) \mid x \in r\}$$

Let x be an element of the radical r such that $\text{ad}(x)$ is an E -bimodule morphism. The proof of Corollary 2.9 shows that x is a linear combination of oriented cycles of length ≥ 1 , so $x = \sum_{(e,\gamma) \in Q_0//B-Q_0} \lambda_\gamma \gamma$ with $\lambda_\gamma \in k$. Since $\xi(\text{ad}(x)) = \sum_{(e,\gamma) \in Q_0//B-Q_0} \lambda_\gamma (\sum_{\substack{a \in eQ_1 \\ \gamma a \in B}} (a, \gamma a) - \sum_{\substack{a \in Q_1 e \\ a\gamma \in B}} (a, a\gamma))$, we are done.

(iv) : It is obvious that the Lie algebra of $\hat{E}$ is generated by the inner derivations $\text{ad}(e)$ where $e \in Q_0$. Since $\xi(\text{ad}(e)) = \sum_{a \in Q_1 e} (a, a) - \sum_{a \in eQ_1} (a, a)$ the proof is finished.

q.e.d. 3.4

Using the notations which will be introduced at the beginning of the next section we get:

Corollary 3.5 *Let k be a field of characteristic 0 and Λ a finite dimensional monomial k -algebra. The Lie algebra of the closed connected normal subgroup $\frac{H_\Lambda \cap U_\Lambda}{H_\Lambda \cap \text{Inn}^*(\Lambda)}$ of the connected algebraic group $\text{Out}(\Lambda)^\circ$ is the Lie ideal $\oplus_{i \geq 1} L_i$ of $H^1(\Lambda, \Lambda)$ and the Lie algebra of the closed connected subgroup $\frac{V_\Lambda^\circ}{E}$ of $\text{Out}(\Lambda)^\circ$ is the Lie subalgebra L_0 of $H^1(\Lambda, \Lambda)$.*

4 The Lie algebra $H^1(\Lambda, \Lambda)$ of a monomial algebra

Since Hochschild cohomology is additive and since its Lie algebra structure follows this additive decomposition we will assume henceforth that the quiver Q is connected. For the study of the Lie algebra $H^1(\Lambda, \Lambda)$ of the monomial algebra $\Lambda = kQ/\langle Z \rangle$ we will use the description which we obtained in Theorem 2.7. For every element $x \in \text{Ker } \psi_1$ we will also denote its class in $H^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$ by x .

If $(a, \gamma) \in Q_1//B_n$ and $(b, \varepsilon) \in Q_1//B_m$, the formula we have obtained shows that $[(a, \gamma), (b, \varepsilon)]$ is an element of $k(Q_1//B_{n+m-1})$. Thus, we have a graduation on the Lie algebra $k(Q_1//B) = \oplus_{i \in \mathbb{N}} k(Q_1//B_i)$ by considering that the elements of $k(Q_1//B_i)$ have degree $i-1$ for all $i \in \mathbb{N}$. It is clear that the Lie subalgebra $\text{Ker } \psi_1$

of $k(Q_1//B)$ preserves this graduation and that $\text{Im } \psi_0$ is a graded ideal. Therefore, the Lie algebra $H^1(\Lambda, \Lambda) = \text{Ker } \psi_1 / \text{Im } \psi_0$ has also a graduation. If we set

$$L_{-1} := k(Q_1//Q_0) \cap \text{Ker } \psi_1$$

$$L_0 := k(Q_1//Q_1) \cap \text{Ker } \psi_1 / \langle \sum_{a \in Q_1 e} (a, a) - \sum_{a \in e Q_1} (a, a) \mid e \in Q_0 \rangle \quad \text{and}$$

$$L_i := k(Q_1//B_{i+1}) \cap \text{Ker } \psi_1 / \langle \sum_{\substack{a \in Q_1 e \\ \gamma a \in B}} (a, \gamma a) - \sum_{\substack{a \in e Q_1 \\ a \gamma \in B}} (a, a \gamma) \mid (e, \gamma) \in Q_0//Q_i \rangle$$

for all $i \geq 1$, $i \in \mathbb{N}$, we obtain $H^1(\Lambda, \Lambda) = \oplus_{i \geq -1} L_i$ and $[L_i, L_j] \subset L_{i+j}$ for all $i, j \geq -1$ where $L_{-2} = 0$.

Lemma 4.1 *L_{-1} equals 0 if and only if there exists for every loop $(a, e) \in Q_1//Q_0$ a path p in Z such that $p^{(a,e)} \neq 0$.*

Proof: Clear, since $\psi_1(a, e) = \sum_{p \in Z} (p, p^{(a,e)})$ for every loop $(a, e) \in Q_1//Q_0$.

q.e.d. 4.1

We study first the subalgebra L_{-1} of $H^1(\Lambda, \Lambda)$.

Proposition 4.2 *Each of the following conditions implies $L_{-1} = 0$:*

- (i) *The quiver Q does not have a loop.*
- (ii) *For every loop $(a, e) \in Q_1//Q_0$ of Q the characteristic of the field k does not divide the integer $m \geq 2$ for which $a^m \in Z$ and $a^{m-1} \in B$.*
- (iii) *The characteristic of the field k is equal to 0.*
- (iv) *Λ is a truncated quiver algebra $kQ/\langle Q_m \rangle$ for a quiver Q different from the loop and for $m \geq 2$.*
- (v) *Q is the loop quiver and $\Lambda = kQ/\langle Q_m \rangle$ is a truncated quiver algebra such that the characteristic of k does not divide $m \geq 2$.*

Proof: (i) : clear

(ii) : Let $(a, e) \in Q_1//Q_0$ be a loop of Q . Since Λ is finite dimensional, there exists an integer $m \geq 2$ such that $p := a^m \in Z$ and $a^{m-1} \in B$. If the characteristic of k does not divide m , then $p^{(a,e)} = ma^{m-1}$ is different from 0 and so $ma^{m-1} \notin \langle Z \rangle$ i.e. $p^{(a,e)} \neq 0$.

(iii) : Clear, because (iii) implies (ii).

(iv) : Using (ii) we can suppose that $\text{char } k$ divides m . Let $(a, e) \in Q_1//Q_0$ be a loop of the quiver Q . By assumption the connected quiver itself is not a loop, therefore there exists an arrow $b \in Q_1$ different from a such that ba^{m-1} or $a^{m-1}b \in Z = Q_m$. The fact that $\text{char } k$ divides m implies that the characteristic of the field k does not divide $m-1$ and thus $0 \neq (m-1)ba^{m-2} \notin \langle Z \rangle = \langle Q_m \rangle$ or $0 \neq (m-1)a^{m-2}b \notin \langle Z \rangle = \langle Q_m \rangle$.

(v) : Clear, because (v) implies (ii).

q.e.d. 4.2

Before we study the case where L_{-1} equals zero we consider the following exceptional case:

Proposition 4.3 *Let Q be the loop having a vertex e and an arrow a . Let $Z = \{a^m\}$ be such that the characteristic p of the field k divides the integer $m \geq 2$ and let $\Lambda = kQ/\langle Z \rangle$ be the monomial algebra associated to Z . The following conditions are equivalent:*

- (i) *The Lie algebra $H^1(\Lambda, \Lambda)$ is simple.*
- (ii) *The Lie algebra $H^1(\Lambda, \Lambda)$ is semisimple.*
- (iii) *The integer $m \geq 2$ is equal to the characteristic p of the field k and $p > 2$.*
- (iv) *$H^1(\Lambda, \Lambda)$ is isomorphic to the Witt Lie algebra $W(1, 1) := \text{Der}(k[X]/(X^p))$ and $p > 2$.*

Proof: (i) $\Rightarrow$ (ii) : trivial

(ii) $\Rightarrow$ (iii) : The assumption $\text{char } k = p$ implies $[L_{-1}, L_{p-1}] = 0$ and thus, $L_{p-1} \oplus \cdots \oplus L_{m-2}$ is a solvable Lie ideal of the Lie algebra $H^1(\Lambda, \Lambda) = \bigoplus_{i=-1}^{m-2} L_i$. Since $H^1(\Lambda, \Lambda)$ is semisimple, we deduce $m = p$. We have $p > 2$, because there is no semisimple Lie algebra of dimension 2.

(iii) $\Rightarrow$ (iv) : Since Λ is isomorphic to the commutative algebra $k[X]/(X^m)$, the Lie algebra $H^1(\Lambda, \Lambda)$ is isomorphic to the Lie algebra of derivations $\text{Der}(k[X]/(X^m))$.

(iv) $\Rightarrow$ (i) : If $\text{char } k > 2$, then the Witt Lie algebra $W(1, 1) := \text{Der}(k[X]/(X^m))$ is one of the non classical simple Lie algebras.

q.e.d. 4.3

Proposition 4.2 shows that the case where L_{-1} is different from zero is quite exceptional. We will assume henceforth that $L_{-1} = 0$. In that case $\bigoplus_{i \geq 1} L_i$ is a solvable Lie ideal of $H^1(\Lambda, \Lambda)$ since $H^1(\Lambda, \Lambda)$ is finite dimensional. It is obvious that L_0 is a Lie subalgebra of $H^1(\Lambda, \Lambda)$ whose bracket is

$$[(a, c), (b, d)] = \delta_{a,d}(b, c) - \delta_{b,c}(a, d)$$

for all $(a, c), (b, d) \in L_0$. It follows that we have $\text{Rad } H^1(\Lambda, \Lambda) = \text{Rad } L_0 \oplus \bigoplus_{i \geq 1} L_i$ and $H^1(\Lambda, \Lambda)/\text{Rad } H^1(\Lambda, \Lambda) = L_0/\text{Rad } L_0$ where $\text{Rad } H^1(\Lambda, \Lambda)$ (resp. $\text{Rad } L_0$) denotes the radical of $H^1(\Lambda, \Lambda)$ (resp. L_0). As a consequence the study of the Lie algebra $H^1(\Lambda, \Lambda)$ can be often reduced to the study of the Lie algebra L_0 .

We recall a few definitions introduced by Guil-Asensio and Saorín (see 2.3 in [9] and 25 in [8]) for the convenience of the reader which will be useful in the following.

Definition 4.4 *Let (a, b) be a couple of parallel arrows. We shall say that the ideal $\langle Z \rangle$ of the algebra kQ is (a, b) -saturated, if for every path p of Z we have $p^{(b,a)} = 0$. This is denoted by $a \leq_{\langle Z \rangle} b$. The ideal $\langle Z \rangle$ is called completely saturated if it is (a, b) -saturated for all $(a, b) \in Q_1//Q_1$.*

Remarks 4.5

- (i) *For every class of parallel arrows $\bar{\alpha}$ the relation $\leq_{\langle Z \rangle}$ on $\bar{\alpha}$ is reflexive and transitive.*
- (ii) *For $p \in Z$ and $(a, b) \in Q_1//Q_1$ we have $p^{(b,a)} = 0$ if and only if each term of the sum is zero (see 2.5), i.e. each replacement of one appearance of b in p by a is a path in Z .*

The hereditary algebra where $Z = \emptyset$ and the truncated algebras where $Z = Q_m$, $m \geq 2$, are examples of monomial algebras whose ideal $\langle Z \rangle$ is completely saturated. Locateli studied the Hochschild cohomology of truncated algebras in [14] and Cibils treated the case of radical square zero algebras (see [2]).

It is clear that the relation of parallelism is an equivalence relation on Q_1 . Denote by $Q_1//$ its set of equivalence classes. We shall call a class of parallel arrows non trivial if it contains at least two arrows. It is easy to check that we have on every class of parallel arrows $\bar{\alpha} = \{\alpha_1, \dots, \alpha_n\} \in Q_1//$ the following equivalence relation

$$\alpha_i \approx \alpha_j \quad := \quad \alpha_i \leq_{\langle Z \rangle} \alpha_j \text{ and } \alpha_j \leq_{\langle Z \rangle} \alpha_i$$

We denote by $\bar{\alpha}/\approx$ its set of equivalence classes. For $R, S \in \bar{\alpha}/\approx$ we write $R \leq_{\langle Z \rangle} S$ if there exist arrows $a \in R$ and $b \in S$ such that $a \leq_{\langle Z \rangle} b$ which is equivalent to saying that $a \leq_{\langle Z \rangle} b$ for all arrows $a \in R$ and $b \in S$. Note that the relation $\leq_{\langle Z \rangle}$ on $\bar{\alpha}/\approx$ is an order relation.

The notion of saturation can be reformulated in terms of the Lie algebra structure:

Remark 4.6 *Let (a, b) be a couple of parallel arrows. The ideal $\langle Z \rangle$ is (a, b) -saturated if and only if (b, a) is an element of the Lie algebra L_0 . Thus the parallel arrows a and b are equivalent if and only if $(a, b) \in L_0$ and $(b, a) \in L_0$. The ideal $\langle Z \rangle$ is completely saturated if and only if every two parallel arrows are equivalent.*

Proposition 4.7

A basis $\mathcal{B}$ of the Lie algebra L_0 is given by the union of the following sets:

- (i) *all the couples $(a, b) \in L_0$ such that the parallel arrows a and b are different*
- (ii) *for every class of parallel arrows $\bar{\alpha} = \{\alpha_1, \dots, \alpha_n\} \in Q_1//$, all the elements $(\alpha_i, \alpha_i) \in L_0$ such that $i < n$*
- (iii) *$|Q_1//| - |Q_0| + 1$ linearly independent elements $(c, c) \in L_0$ different from those in (ii).*

Proof: An element $\sum_{(a,b) \in Q_1//Q_1} \lambda_{(a,b)}(a, b) \in k(Q_1//Q_1)$, with $\lambda_{(a,b)} \in k$, is contained in $\text{Ker } \psi_1$ if and only if

$$0 = \sum_{(a,b) \in Q_1//Q_1} \lambda_{(a,b)} \psi_1(a, b) = \sum_{\substack{(a,b) \in Q_1//Q_1 \\ a \neq b}} \lambda_{(a,b)} \sum_{p \in Z} (p, p^{(a,b)})$$

First we recall that for a couple of distinct arrows (a, b) , the non zero terms of the sum defining $p^{(a,b)}$ are all distinct (see 2.5), forming a subset of the basis $Z//B$. Second we check the following: if $a \neq b$ and if $(p, p^{(a,b)}) = (q, q^{(c,d)})$ then $p = q$ and if $p^{(a,b)} \neq 0$ then $(a, b) = (c, d)$. Indeed let $p = p_n \dots p_2 p_1$ be a path of Z of length l and consider the summand $p_n \dots p_{i+1} b p_{i-1} \dots p_1 \in B$ of $p^{(a,b)}$ obtained by replacing the arrow $p_i = a$ by $b \neq a$. The fact $(p, p^{(a,b)}) = (p, p^{(c,d)}) \neq 0$ implies that there exists $j \in \{1, \dots, n\}$ such that $p_n \dots p_{j+1} c p_{j-1} \dots p_{i+1} b p_{i-1} \dots p_1 = p_n \dots p_{j+1} d p_{j-1} \dots p_{i+1} a p_{i-1} \dots p_1$. Since $a \neq b$, we have $i = j$ as well as $a = d$ and $c = b$. This shows that if $a \neq b$ then the non zero elements $(p, p^{(a,b)})$ of $k(Z//B)$ are linearly independent. Therefore we can choose a basis of $k(Q_1//Q_1) \cap \text{Ker } \psi_1$ in $Q_1//Q_1$: take all elements $(a, a) \in Q_1//Q_1$ and all elements $(a, b) \in Q_1//Q_1$, $a \neq b$, such that the ideal $\langle Z \rangle$ is (b, a) -saturated. We have for every vertex $e \in Q_0$ in $L_0 = k(Q_1//Q_1) \cap \text{Ker } \psi_1 / k(Q_1//Q_1) \cap \text{Im } \psi_0$ the relation $\sum_{a \in Q_1 e} (a, a) - \sum_{a \in e Q_1} (a, a) = 0$. Since $|Q_0| - 1$ of those $|Q_0|$ relations are linearly independent, we see that $\dim k(Q_1//Q_1) \cap \text{Im } \psi_0 = |Q_1| - |Q_0| + 1$. Thus we get a basis of L_0 if we impose the conditions (ii) and (iii).

q.e.d. 4.7

Let $\mathcal{B}$ be a basis of L_0 as described in the preceding Proposition.

Remark 4.8 *If the quiver Q has a (non oriented or oriented) cycle, then there exists at least one class of parallel arrows $\overline{\alpha} = \{\alpha_1, \dots, \alpha_n\}$, $n \geq 1$, such that $\mathcal{B}$ contains all the elements $(\alpha_i, \alpha_i) \in L_0$, $1 \leq i \leq n$.*

For every class of parallel arrows $\overline{\alpha} \in Q_1//$ we denote by $L_0^{\overline{\alpha}}$ the Lie ideal of L_0 generated by the elements $(a, b) \in \mathcal{B}$ such that $a, b \in \overline{\alpha}$. Obviously the Lie algebra L_0 is the product of these Lie algebras:

$$L_0 = \prod_{\overline{\alpha} \in Q_1//} L_0^{\overline{\alpha}}$$

To study the radical of this Lie algebra we need the following Lemma:

Lemma 4.9 *Let $J \neq 0$ be a Lie ideal of $L_0^{\overline{\alpha}}$ generated by elements $(\alpha_i, \alpha_j) \in L_0^{\overline{\alpha}}$ such that $(\alpha_j, \alpha_i) \notin L_0^{\overline{\alpha}}$. Then $\dim_k[J, J] < \dim_k J$.*

Proof: Let (α_i, α_j) be any element of J . If $(\alpha_i, \alpha_j) \notin [J, J]$ then nothing is to show. So suppose $(\alpha_i, \alpha_j) \in [J, J]$. This is the case if and only if there exists an arrow $\alpha_l \in \overline{\alpha}$, $\alpha_i \neq \alpha_l \neq \alpha_j$, such that $(\alpha_i, \alpha_l) \in J$ and $(\alpha_l, \alpha_j) \in J$. If $(\alpha_i, \alpha_l) \notin [J, J]$ then we are done. If not we start again. Suppose that we have $(\alpha_i, \alpha_l) \in [J, J]$. Thus there exists an arrow $\alpha_k \in \overline{\alpha} - \{\alpha_i, \alpha_l\}$ such that $(\alpha_i, \alpha_k) \in J$ and $(\alpha_k, \alpha_l) \in J$. The fact $(\alpha_l, \alpha_j) \in J$ implies $(\alpha_j, \alpha_l) \notin J$ and so $\alpha_k \neq \alpha_j$. If $(\alpha_i, \alpha_k) \notin [J, J]$ then we are done. If not we start again. Since $\overline{\alpha}$ is a finite set, this procedure stops after a finite number of steps.

q.e.d. 4.9

Theorem 4.10 *The radical of the Lie algebra L_0 is generated as a k -vector space by the following elements of L_0 : for every class of parallel arrows $\overline{\alpha} = \{\alpha_1, \dots, \alpha_n\}$*

- $\sum_{\alpha_i \in S} (\alpha_i, \alpha_i)$ for every equivalence class $S \in \overline{\alpha}/\approx$
- (α_i, α_j) , $i \neq j$, if $(\alpha_i, \alpha_j) \in L_0$ and $(\alpha_j, \alpha_i) \notin L_0$
- $(\alpha_{i_1}, \alpha_{i_2})$, $(\alpha_{i_2}, \alpha_{i_1})$ and $(\alpha_{i_1}, \alpha_{i_1})$ for all $S = \{\alpha_{i_1}, \alpha_{i_2}\} \in \overline{\alpha}/\approx$ if $\text{char } k = 2$

Proof: Let I be the k -vector space generated by the above-described elements. If we define $I_{\overline{\alpha}} := I \cap L_0^{\overline{\alpha}}$ we obtain $I = \prod_{\overline{\alpha} \in Q_1//} I_{\overline{\alpha}}$. Let $\overline{\alpha} = \{\alpha_1, \dots, \alpha_n\}$ be a class of parallel arrows of Q . It is easy to check using the definition of the bracket on L_0 , that $I_{\overline{\alpha}}$ is a Lie ideal of $L_0^{\overline{\alpha}}$ and that $I_{\overline{\alpha}}^{(3)} := [I_{\overline{\alpha}}^{(2)}, I_{\overline{\alpha}}^{(2)}]$ only contains elements $(\alpha_i, \alpha_j) \in L_0^{\overline{\alpha}}$ such that $(\alpha_i, \alpha_j) \notin L_0^{\overline{\alpha}}$. From the fact that $\dim_k I_{\overline{\alpha}} < \infty$ we deduce successively using the preceding Lemma that there exists an $l \in \mathbb{N}$ such that $I_{\overline{\alpha}}^{(l)} := [I_{\overline{\alpha}}^{(l-1)}, I_{\overline{\alpha}}^{(l-1)}] = 0$. Thus $I_{\overline{\alpha}}$ is a solvable Lie ideal of $L_0^{\overline{\alpha}}$ and we are done if we show that $L_0^{\overline{\alpha}}/I_{\overline{\alpha}}$ is a semisimple Lie algebra or 0. From the relations in $L_0^{\overline{\alpha}}/I_{\overline{\alpha}}$ follows that we obtain a basis of $L_0^{\overline{\alpha}}/I_{\overline{\alpha}}$ by taking all elements (α_i, α_j) , where $\alpha_i \neq \alpha_j$ and $\alpha_i, \alpha_j \in S$, and $|S| - 1$ elements (α_i, α_i) where $\alpha_i \in S$ for every $S \in \overline{\alpha}/\approx$, with $|S| > 2$ in case $\text{char } k = 2$. For every $S \in \overline{\alpha}/\approx$ we denote by ${}^S L_0^{\overline{\alpha}}/I_{\overline{\alpha}}$ the Lie ideal generated by the elements $(\alpha_i, \alpha_j) \in L_0^{\overline{\alpha}}$ such that $\alpha_i, \alpha_j \in S$. Then we have $L_0^{\overline{\alpha}}/I_{\overline{\alpha}} = \prod_{S \in \overline{\alpha}/\approx} {}^S L_0^{\overline{\alpha}}/I_{\overline{\alpha}}$. Let $S = \{\alpha_{i_1}, \dots, \alpha_{i_m}\}$ be an equivalence class in $\overline{\alpha}/\approx$, with $m \geq 2$ if $\text{char } k \neq 2$ and $m \neq 2$ if $\text{char } k = 2$. The elements $(\alpha_{i_p}, \alpha_{i_q})$, $i_p \neq i_q$, and $(\alpha_{i_p}, \alpha_{i_p})$, $i_p \neq i_m$, form a basis of ${}^S L_0^{\overline{\alpha}}/I_{\overline{\alpha}}$ and we have the relation

$\sum_{\alpha_{i_p} \in S} (\alpha_{i_p}, \alpha_{i_p}) = 0$. As the bracket on ${}^S L_0^{\bar{\alpha}} / I_{\bar{\alpha}}$ is given by $[(\alpha_{i_p}, \alpha_{i_q}), (\alpha_{i_k}, \alpha_{i_l})] = \delta_{i_q, i_k} (\alpha_{i_p}, \alpha_{i_l}) - \delta_{i_p, i_l} (\alpha_{i_k}, \alpha_{i_q})$ we see that ${}^S L_0^{\bar{\alpha}} / I_{\bar{\alpha}} \rightarrow \mathfrak{pgl}(m, k) := \mathfrak{gl}(m, k) / k1$ given by $(\alpha_{i_p}, \alpha_{i_q}) \mapsto e_{pq} \bmod k1$ is a Lie algebra isomorphism. Since $\mathfrak{pgl}(m, k)$ is a semisimple Lie algebra if $m \geq 2$ when $\text{char } k \neq 2$ and if $m > 2$ when $\text{char } k = 2$, we conclude that $L_0^{\bar{\alpha}} / I_{\bar{\alpha}}$ is a semisimple Lie algebra.

q.e.d. 4.10

Corollary 4.11 *The semisimple Lie algebra $H^1(\Lambda, \Lambda) / \text{Rad } H^1(\Lambda, \Lambda) = L_0 / \text{Rad } L_0$ is the product of Lie algebras having a factor $\mathfrak{pgl}(|S|, k) := \mathfrak{gl}(|S|, k) / k1$ for every equivalence class $S \in \bar{\alpha} / \approx$ of a class of parallel arrows $\bar{\alpha}$ such that $|S| \geq 2$ if $\text{char } k \neq 2$ and $|S| > 2$ if $\text{char } k = 2$.*

Corollary 4.12 *Let Q be a connected quiver and $\Lambda = kQ / \langle Z \rangle$ a finite dimensional monomial algebra such that $L_{-1} = k(Q_0 / Q_1) \cap \text{Ker } \psi_1 = 0$. The following conditions are equivalent:*

- (i) *The Lie algebra $H^1(\Lambda, \Lambda)$ is solvable.*
- (ii) *Every equivalence class S of a class of parallel arrows $\bar{\alpha} \in Q_1 / /$ of Q contains one and only one arrow if the characteristic of the field k is not 2. In the case $\text{char } k = 2$ we have $|S| \leq 2$ for all $S \in \bar{\alpha} / \approx$, $\bar{\alpha} \in Q_1 / /$.*

Since in characteristic 0 the Lie algebra of a connected algebraic group is solvable if and only if the connected algebraic group is solvable (as a group), using Proposition 4.2 we infer the following result also proved by Guil-Asensio and Saorín (see Corollary 2.22 in [9]):

Corollary 4.13 *Let k be a field of characteristic 0, Q a connected quiver and $\Lambda = kQ / \langle Z \rangle$ a finite dimensional monomial algebra. Then the identity component of the algebraic group of outer automorphisms $\text{Out}(\Lambda)^\circ$ is solvable if and only if every two parallel arrows of Q are not equivalent.*

Definition 4.14 *Let $\bar{Q}$ be the subquiver of Q obtained by taking a representative for every class of parallel arrows of Q .*

Theorem 4.15 *Let Q be a connected quiver and $\Lambda = kQ / \langle Z \rangle$ a finite dimensional monomial algebra such that $L_{-1} = k(Q_0 / Q_1) \cap \text{Ker } \psi_1 = 0$. The following conditions are equivalent:*

- (i) *The Lie algebra $H^1(\Lambda, \Lambda)$ is semisimple.*
- (ii) *The quiver $\bar{Q}$ is a tree, Q has at least one non trivial class of parallel arrows and the ideal $\langle Z \rangle$ is completely saturated. If the characteristic of the field k is 2, then Q does not have a class of parallel arrows containing exactly two arrows.*
- (iii) *$H^1(\Lambda, \Lambda)$ is isomorphic to the non trivial product of Lie algebras*

$$\prod_{\bar{\alpha} \in Q_1 / /} \mathfrak{pgl}(|\bar{\alpha}|, k)$$

where $|\bar{\alpha}| \neq 2$ if the characteristic of k is equal to 2.

Proof: Let $\mathcal{B}$ be a basis of L_0 as described in Proposition 4.7.

(i) $\Rightarrow$ (ii) : If $\overline{Q}$ contained a (non oriented or oriented) cycle, there would exist a class $\overline{\alpha} \in Q_1//$ of parallel arrows such that $(\alpha, \alpha) \in \mathcal{B}$ for all $\alpha \in \overline{\alpha}$ (see 4.8) and so $k(\sum_{\alpha \in \overline{\alpha}} (\alpha, \alpha))$ would be a nontrivial abelian ideal of L_0 which contradicts the fact that $0 = \text{Rad } H^1(\Lambda, \Lambda) = \text{Rad } L_0 \oplus \bigoplus_{i \geq 1} L_i$. To insure that $H^1(\Lambda, \Lambda) = L_0 \neq 0$ it is necessary that Q contains at least one nontrivial class of parallel arrows. Suppose that $\langle Z \rangle$ is not completely saturated. According to Remark 4.6 there exist two parallel arrows which are not equivalent. This means that there exists a class of parallel arrows $\overline{\alpha} \in Q_1//$ such that $\overline{\alpha}/\approx$ contains at least two elements, say R and S . The properties of the basis $\mathcal{B}$ show that either $\sum_{a \in R} (a, a) \neq 0$ or $\sum_{a \in S} (a, a) \neq 0$ which contradicts the fact that $\text{Rad } L_0 \subset \text{Rad } H^1(\Lambda, \Lambda) = 0$. The last statement is an immediate consequence of Theorem 4.10.

(ii) $\Rightarrow$ (iii) : Let Q and Z satisfy the given conditions. From the fact that $\overline{Q}$ is a tree, we deduce $\bigoplus_{i \geq 1} L_i = 0$ and so $H^1(\Lambda, \Lambda) = L_0$. Since $\langle Z \rangle$ is completely saturated we have $L_0 = k(Q_1//Q_1)/\langle \sum_{a \in Q_1 e} (a, a) - \sum_{a \in e Q_1} (a, a) \mid e \in Q_0 \rangle$. This and the fact that Q does not have an equivalence class containing two arrows if $\text{char } k = 2$, show that its radical is generated by the elements of type $\sum_{\alpha \in \overline{\alpha}} (\alpha, \alpha)$, $\overline{\alpha} \in Q_1//$, (see Theorem 4.10). These elements being equal to zero we see that $\text{Rad } H^1(\Lambda, \Lambda) = 0$ and Corollary 4.11 yields the result.

(iii) $\Rightarrow$ (i) : Clear in view of the following Remark.

q.e.d. 4.15

Remark 4.16 *The Lie algebra $\mathfrak{pgl}(n, k) := \mathfrak{gl}(n, k)/k1$, $n \geq 2$, is isomorphic to the classical simple Lie algebra $\mathfrak{sl}(n, k)$ of $n \times n$ -matrices having trace zero if the characteristic of the field k does not divide n . If $\text{char } k$ divides n and $n \neq 2$, then $\mathfrak{pgl}(n, k)$ is a semisimple algebra without being a direct product of simple Lie algebras.*

Corollary 4.17 *Let Q be a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra such that $L_{-1} = k(Q_0//Q_1) \cap \text{Ker } \psi_1 = 0$. The following conditions are equivalent:*

- (i) *The Lie algebra $H^1(\Lambda, \Lambda)$ is simple.*
- (ii) *The quiver $\overline{Q}$ is a tree, the quiver Q has exactly one class of parallel arrows $\overline{\alpha} = \{\alpha_1, \dots, \alpha_n\}$ such that $n \geq 2$ and the characteristic of the field k does not divide n . The ideal $\langle Z \rangle$ is completely saturated.*
- (iii) *There exists an integer $n \geq 2$ such that the characteristic of the field k does not divide n and $H^1(\Lambda, \Lambda)$ is isomorphic to the Lie algebra $\mathfrak{sl}(n, k)$.*

Since the first Hochschild cohomology group is the Lie algebra of the algebraic group of outer automorphisms in characteristic 0 (see Proposition 3.1), we get the following Corollary (see Corollary 4.9 in [9] for a partial result on hereditary algebras):

Corollary 4.18 *Let k be a field of characteristic 0, Q a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra. The following conditions are equivalent:*

- (i) *The algebraic group $\text{Out } (\Lambda)^\circ$ is semisimple.*
- (ii) *The quiver $\overline{Q}$ is a tree, the quiver Q has at least one non trivial class of parallel arrows and the ideal $\langle Z \rangle$ is completely saturated.*

(iii) The algebraic group $\text{Out}(\Lambda)^\circ$ is isomorphic to the non trivial product of algebraic groups $\prod_{\bar{\alpha} \in Q_1//} \mathbf{PGL}(|\bar{\alpha}|, k)$.

Proof: In characteristic 0 a connected algebraic group is semisimple if and only if its Lie algebra is semisimple (see [12] 13.5).

(i) $\Leftrightarrow$ (ii) : This equivalence follows immediately from the preceding Theorem and Proposition 3.1.

(ii) $\Rightarrow$ (iii) : Let Q and Z satisfy the given condition. According to 3.3 we know that $\text{Out}(\Lambda)^\circ = \frac{H_\Lambda \cap U_\Lambda}{H_\Lambda \cap \text{Inn}^*(\Lambda)} \rtimes \frac{V_\Lambda^\circ}{\hat{E}}$. Since $\bar{Q}$ is a tree, there is no path in $B' := B - Q_0 \cup Q_1$ of length ≥ 2 which is parallel to an arrow. From the fact that we have for every element $\sigma \in H_\Lambda \cap U_\Lambda$ that $\sigma(a) = a + \sum_{(a, \gamma) \in Q_1//B'} \lambda_{(a, \gamma)} \gamma$ for all $a \in Q_1$ we deduce that $H_\Lambda \cap U_\Lambda = \{\text{id}_\Lambda\}$. As the ideal $\langle Z \rangle$ is completely saturated by assumption, any two parallel arrows are equivalent. Therefore the main Theorem 2.20 of the article [9] implies $V_\Lambda^\circ \simeq \prod_{\bar{\alpha} \in Q_1//} \mathbf{GL}(|\bar{\alpha}|, k)$. Since the quiver $\bar{Q}$ is assumed to be a tree, we have $\hat{E} = \text{Ch}(Q, k) = \prod_{\bar{\alpha} \in Q_1//} k^* I_{|\bar{\alpha}|}$ thanks to example 8 in [8]. Hence we obtain

$$\begin{aligned} \text{Out}(\Lambda)^\circ &= \frac{H_\Lambda \cap U_\Lambda}{H_\Lambda \cap \text{Inn}^*(\Lambda)} \rtimes \frac{V_\Lambda^\circ}{\hat{E}} = \frac{V_\Lambda^\circ}{\hat{E}} \\ &\simeq \frac{\prod_{\bar{\alpha} \in Q_1//} \mathbf{GL}(|\bar{\alpha}|, k)}{\prod_{\bar{\alpha} \in Q_1//} k^* I_{|\bar{\alpha}|}} = \prod_{\bar{\alpha} \in Q_1//} \frac{\mathbf{GL}(|\bar{\alpha}|, k)}{k^* I_{|\bar{\alpha}|}} = \prod_{\bar{\alpha} \in Q_1//} \mathbf{PGL}(|\bar{\alpha}|, k) \end{aligned}$$

(iii) $\Rightarrow$ (i) : The semisimplicity of the Lie algebra $\mathcal{L}(\text{Out}(\Lambda)^\circ) = \prod_{\bar{\alpha} \in Q_1//} \mathfrak{pgl}(|\bar{\alpha}|, k)$ implies the semisimplicity of the connected algebraic group $\text{Out}(\Lambda)^\circ$.

q.e.d. 4.18

The fact that in characteristic 0 the Lie algebra of a connected algebraic group is simple if and only if the connected algebraic group is almost simple, yields the following result:

Corollary 4.19 *Let k be a field of characteristic 0, Q a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra. The following conditions are equivalent:*

- (i) *The algebraic group $\text{Out}(\Lambda)^\circ$ is almost simple.*
- (ii) *The quiver $\bar{Q}$ is a tree, the quiver Q has exactly one class of parallel arrows $\bar{\alpha} = \{\alpha_1, \dots, \alpha_n\}$ of order $n \geq 2$ and the ideal $\langle Z \rangle$ is completely saturated.*
- (iii) *There exists an integer $n \geq 2$ such that the algebraic group $\text{Out}(\Lambda)^\circ$ is isomorphic to the algebraic group $\mathbf{PGL}(n, k)$.*

In order to get criteria for the commutativity and the reductivity of the Lie algebra $H^1(\Lambda, \Lambda)$ we need to study its center. Therefore we introduce the following definitions: for every class of parallel arrows $\bar{\alpha}$ of Q we call a set $C \subset \bar{\alpha}$ connected, if for every two arrows α_1 and α_r of C there exist arrows $\alpha_2, \dots, \alpha_{r-1} \in C$ such that we have $\alpha_i \leq_{\langle Z \rangle} \alpha_{i+1}$ or $\alpha_{i+1} \leq_{\langle Z \rangle} \alpha_i$ for all $i \in \{1, \dots, r-1\}$. A connected set $C \subset \bar{\alpha}$ is called a connected component of $\bar{\alpha}$ if it is maximal for the connection, i.e. for every arrow $\beta \in \bar{\alpha} - C$ there is no arrow $\alpha \in C$ such that $\alpha \leq_{\langle Z \rangle} \beta$ or $\beta \geq_{\langle Z \rangle} \alpha$. Clearly the connected components of a class of parallel arrows $\bar{\alpha}$ form a partition of $\bar{\alpha}$.

Lemma 4.20 *Let Q be a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra.*

- (i) *The center $Z(L_0)$ of the Lie algebra L_0 is generated by the elements $\sum_{a \in C} (a, a)$ where C denotes a connected component of a class of parallel arrows of Q .*

(ii) If the field k has characteristic 0 or if the quiver Q does not have an oriented cycle, then the center $Z(H^1(\Lambda, \Lambda))$ of the Lie algebra $H^1(\Lambda, \Lambda)$ is contained in the center $Z(L_0)$ of L_0 .

Proof: (i) : For every element $(a, b) \in Q_1//Q_1$, $a \neq b$, we have $[(a, b), (a, a)] = (a, b)$ so that $Z(L_0)$ is contained in the abelian Lie subalgebra of L_0 generated by the elements (a, a) , $a \in Q_1$. For every linear combination $\sum_{a \in Q_1} \lambda_a (a, a)$, $\lambda_a \in k$, we have $[\sum_{a \in Q_1} \lambda_a (a, a), (b, c)] = (-\lambda_b + \lambda_c)(b, c)$ for all $(b, c) \in L_0$. Therefore $\sum_{a \in Q_1} \lambda_a (a, a)$ is contained in $Z(L_0)$ if and only if we have $\lambda_b = \lambda_c$ for all arrows b, c such that $c \geq_{\langle Z \rangle} b$. This shows that $Z(L_0)$ is generated by the elements $\sum_{a \in C} (a, a)$. (ii) : In both cases we have $L_{-1} = 0$ according to Proposition 4.2. If the characteristic of k is 0, then we have for every element $0 \neq \sum_{(a, \gamma) \in Q_1//B_{i+1}} \lambda_{(a, \gamma)} (a, \gamma) \in L_i$, $\lambda_{(a, \gamma)} \in k$, $i \geq 1$, that

$$\left[\sum_{(a, \gamma) \in Q_1//B_{i+1}} \lambda_{(a, \gamma)} (a, \gamma), \sum_{b \in Q_1} (b, b) \right] = -i \sum_{(a, \gamma) \in Q_1//B_{i+1}} \lambda_{(a, \gamma)} (a, \gamma) \neq 0$$

This shows $Z(H^1(\Lambda, \Lambda)) \subset Z(L_0)$. We assume now that Q does not have an oriented cycle. Then every path $\gamma \in B$ of length ≥ 2 parallel to an arrow a cannot contain a and so $[(a, \gamma), (a, a)] = (a, \gamma)$. Taking into account the fact $H^1(\Lambda, \Lambda) = L_0 \oplus \bigoplus_{i \geq 1} k(Q_1//B_{i+1}) \cap \text{Ker } \psi_1$ yields $Z(H^1(\Lambda, \Lambda)) \subset Z(L_0)$.

q.e.d. 4.20

Recall that a Lie algebra is called reductive if its radical and its center are equal.

Proposition 4.21 *Let Q be a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra. If Q does not have an oriented cycle or if the field k has characteristic 0, then the following conditions are equivalent:*

- (i) *The Lie algebra $H^1(\Lambda, \Lambda)$ is reductive.*
- (ii) *The Lie ideal $\bigoplus_{i \geq 1} L_i$ is equal to 0 and the relation $\leq_{\langle Z \rangle}$ is symmetric on every class of parallel arrows. If the characteristic of k is 2, then there exists no equivalence class $S \in \bar{\alpha}/\approx$ containing exactly two arrows of a class of parallel arrows $\bar{\alpha}$.*

Proof: (i) $\Rightarrow$ (ii) : Let be $\text{Rad } H^1(\Lambda, \Lambda) = Z(H^1(\Lambda, \Lambda))$. If Q does not have an oriented cycle or if $\text{char } k = 0$, then $L_{-1} = 0$ and $\text{Rad } H^1(\Lambda, \Lambda) = \text{Rad } L_0 \oplus \bigoplus_{i \geq 1} L_i$. The preceding Lemma shows that the center of $H^1(\Lambda, \Lambda)$ is included in L_0 and thus $\text{Rad } H^1(\Lambda, \Lambda) \subset L_0$. This implies $\bigoplus_{i \geq 1} L_i = 0$ and therefore $H^1(\Lambda, \Lambda) = L_0$. In view of Theorem 4.10 and the preceding Lemma, the assumption $\text{Rad } L_0 = \text{Rad } H^1(\Lambda, \Lambda) = Z(H^1(\Lambda, \Lambda)) = Z(L_0)$ implies that there is no element $(a, b) \in L_0$ such that $(b, a) \notin L_0$, i.e. the relation $\leq_{\langle Z \rangle}$ is symmetric. Furthermore there exists no equivalence class $S \in \bar{\alpha}/\approx$, $\bar{\alpha} \in Q_1//$, containing exactly two elements if $\text{char } k = 2$.

(ii) $\Rightarrow$ (i) : The assumptions imply $L_{-1} = 0$. From $\bigoplus_{i \geq 1} L_i = 0$ we deduce $H^1(\Lambda, \Lambda) = L_0$. Theorem 4.10 and the preceding Lemma show that $\text{Rad } H^1(\Lambda, \Lambda) = \text{Rad } L_0 = Z(L_0) = Z(H^1(\Lambda, \Lambda))$.

q.e.d. 4.21

Since in characteristic 0 a connected algebraic group is reductive if and only if its Lie algebra is reductive, we deduce immediately using Proposition 3.1 and Corollary 3.5

Corollary 4.22 *Let k be a field of characteristic 0, Q a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra. Then the identity component of the algebraic group of the outer automorphisms $\text{Out}(\Lambda)^\circ$ is reductive if and only if the relation $\leq_{\langle Z \rangle}$ is symmetric on every class of parallel arrows of Q and if $\frac{H_\Lambda \cap U_\Lambda}{H_\Lambda \cap \text{Inn}^*(\Lambda)} = 0$.*

Define a sequence of ideals of a Lie algebra L by setting $\mathcal{C}^0 L := L (= L^{(0)})$, $\mathcal{C}^1 L := [L, L] (= L^{(1)})$, $\mathcal{C}^2 L := [L, \mathcal{C}^1 L]$, $\dots$, $\mathcal{C}^n L := [L, \mathcal{C}^{n-1} L]$. A Lie algebra L is called nilpotent if there exists a nonnegative integer m such that $\mathcal{C}^m L = 0$. The integer m such that $\mathcal{C}^m L = 0$ and $\mathcal{C}^{m-1} L \neq 0$ is called the nilindex of L . A Lie algebra L is called filiform if it is nilpotent of maximal nilindex m that is $\dim \mathcal{C}^k L = m - k$ for $1 \leq k \leq m$. The fact $L^{(n)} \subset \mathcal{C}^n L$ for all $n \in \mathbb{N}$ implies that nilpotent algebras are solvable.

Proposition 4.23 *Let Q be a connected quiver and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra. If Q does not have an oriented cycle or if $\text{char } k = 0$, then the following conditions are equivalent:*

- (i) *The Lie algebra $H^1(\Lambda, \Lambda)$ is abelian.*
- (ii) *The Lie algebra $H^1(\Lambda, \Lambda)$ is filiform.*
- (iii) *The Lie algebra $H^1(\Lambda, \Lambda)$ is nilpotent.*
- (iv) *$\bigoplus_{i \geq 1} L_i = 0$ and there exist no parallel arrows $a \neq b$ satisfying $a \leq_{\langle Z \rangle} b$.*
- (v) *The Lie algebra $H^1(\Lambda, \Lambda)$ is generated by the elements (a, a) of L_0 .*
- (vi) *The dimension of the Lie algebra $H^1(\Lambda, \Lambda)$ equals the Euler characteristic $|Q_1| - |Q_0| - 1$.*

Proof: (i) $\Rightarrow$ (ii) and (ii) $\Rightarrow$ (iii) are obvious.

(iii) $\Rightarrow$ (iv) : If there were two parallel arrows $a \neq b$ such that $a \leq_{\langle Z \rangle} b$, we would have $(b, a) \in L_0$ and $[(a, a), (b, a)] = (b, a)$, and so $(b, a) \in \mathcal{C}^n H^1(\Lambda, \Lambda)$ for all $n \in \mathbb{N}$, contrary to the assumption. If Q does not have an oriented cycle, then there exists for every element $0 \neq \sum_{(a, \gamma) \in Q_1 // B_{i+1}} \lambda_{(a, \gamma)}(a, \gamma) \in L_i$, $i \geq 1$, an arrow $b \in Q_1$ such that $[\sum_{(a, \gamma) \in Q_1 // B_{i+1}} \lambda_{(a, \gamma)}(a, \gamma), (b, b)] = \sum_{(b, \gamma) \in Q_1 // B_{i+1}} \lambda_{(b, \gamma)}(b, \gamma) \neq 0$ and $[\sum_{(b, \gamma) \in Q_1 // B_{i+1}} \lambda_{(b, \gamma)}(b, \gamma), (b, b)] = \sum_{(b, \gamma) \in Q_1 // B_{i+1}} \lambda_{(b, \gamma)}(b, \gamma) \neq 0$. Thus we have $0 \neq \sum_{(b, \gamma) \in Q_1 // B_{i+1}} \lambda_{(b, \gamma)}(b, \gamma) \in \mathcal{C}^n H^1(\Lambda, \Lambda)$ for all $n \in \mathbb{N}$. Since the Lie algebra $H^1(\Lambda, \Lambda)$ is assumed to be nilpotent, it follows that $\bigoplus_{i \geq 1} L_i = 0$. In case $\text{char } k = 0$ we have for every element $\sum_{(a, \gamma) \in Q_1 // B_{i+1}} \lambda_{(a, \gamma)}(a, \gamma) \in L_i$, $i \geq 1$,

$$[\sum_{(a, \gamma) \in Q_1 // B_{i+1}} \lambda_{(a, \gamma)}(a, \gamma), \sum_{b \in Q_1} (b, b)] = -i \sum_{(a, \gamma) \in Q_1 // B_{i+1}} \lambda_{(a, \gamma)}(a, \gamma) \neq 0$$

and thus $\sum_{(a, \gamma) \in Q_1 // B_{i+1}} \lambda_{(a, \gamma)}(a, \gamma) \in \mathcal{C}^n H^1(\Lambda, \Lambda)$ for all $n \in \mathbb{N}$ which implies $\bigoplus_{i \geq 1} L_i = 0$.

(iv) $\Rightarrow$ (v) : By assumption $H^1(\Lambda, \Lambda) = L_0$ and L_0 does not contain an element (a, b) , $a \neq b$.

(v) $\Rightarrow$ (vi) : We deduce from the given condition that $\dim \text{Ker } \psi_1 = |Q_1|$. Since the dimension of $\text{Im } \psi_0 = \langle \sum_{a \in Q_1 e} (a, a) - \sum_{a \in e Q_1} (a, a) \mid e \in Q_0 \rangle$ is equal to $|Q_0| - 1$ we obtain $\dim H^1(\Lambda, \Lambda) = \dim \text{Ker } \psi_1 - \dim \text{Im } \psi_0 = |Q_1| - |Q_0| + 1$.

(vi) $\Rightarrow$ (i) : From $\dim \text{Im } \psi_0 = |Q_0| - 1$ and $\dim H^1(\Lambda, \Lambda) = |Q_1| - |Q_0| + 1$ it follows that $\dim \text{Ker } \psi_1 = |Q_1|$ and thus $\text{Ker } \psi_1$ is generated by the elements $(a, a) \in Q_1 // Q_1$. Since the bracket of the Lie algebra $k(Q_1 // B)$ is such that $[(a, a), (b, b)] = 0$ for all $a, b \in Q_1$, we see that $H^1(\Lambda, \Lambda)$ is an abelian Lie algebra.

The following Corollary is clear thanks to Proposition 3.1, because in characteristic 0 a connected algebraic group is abelian (resp. nilpotent) if and only if its Lie algebra is abelian (resp. nilpotent) (see [12] 13.4 and 10.5). It generalizes Guil-Asensio and Saorín's criterion on the commutativity of the algebraic group $V_\Lambda^\circ (= \varepsilon_\Lambda(G_\Lambda))$ (see Corollary 2.23 in [9]).

Corollary 4.24 *Let k be a field of characteristic 0 and $\Lambda = kQ/\langle Z \rangle$ a finite dimensional monomial algebra. The following conditions are equivalent:*

- (i) *The algebraic group $\text{Out}(\Lambda)^\circ$ is abelian.*
- (ii) *The algebraic group $\text{Out}(\Lambda)^\circ$ is nilpotent.*
- (iii) *For every path $\gamma \in B$ parallel to an arrow $a \neq \gamma$ there exists a path p of Z such that $p^{(a,\gamma)} \neq 0$.*

5 Application to group algebras where the group admits a normal cyclic Sylow p -subgroup

The starting point for the following application is the paragraph ‘Représentations modulaires des groupes finis’ in Gabriel’s article [3] (see also [4] p.75). Let k be an algebraically closed field and G a finite group. If the characteristic of k does not divide the order of G , then Maschke’s Theorem states that the group algebra kG is semisimple. Since k is supposed to be algebraically closed, this is equivalent to saying that the group algebra kG is separable and thus $H^1(kG, kG) = 0$. So henceforth let k be a field of characteristic p dividing the order of G . Let $n = p^a q$ be the order of G with q prime to p . We are interested in this section in the particular case where G contains only one cyclic Sylow p -subgroup S , necessarily normal in G . Then S is a normal Hall subgroup of G and Schur’s splitting theorem (see [16] 9.3.6) states that there exists a supplement K of S in G , i.e. a subgroup such that $S \cap K = \{1\}$ and $SK = G$. It is unique up to a group isomorphism. If σ is a generator of S , the action of K on S by conjugation is given by a formula of the type

$$x\sigma x^{-1} = \sigma^{\chi(x)}$$

with $x \in K$ and $\chi(x) \in (\mathbb{Z}/p^a\mathbb{Z})^\times$. The group G is the semidirect product of S and K . For every kK -module N we denote by ${}_\chi N$ the underlying k -vector space of N equipped with a new action $*$ of K such that we have

$$x * m := \chi(x)x \cdot m$$

for all $x \in K$ and $m \in N$. If the kK -module N belongs to the isomorphism class e , we denote by χe the isomorphism class of the kK -module ${}_\chi N$. That which enables us to study the Lie algebra $H^1(kG, kG)$ is the association of a quiver Q to the group G in the following way: the set Q_0 of vertices of Q consists in the set of isomorphism classes of simple kK -modules and for every vertex e we take an arrow $e \rightarrow \chi e$. The quiver Q is a disjoint union of crowns i.e. of oriented cycles. Gabriel proved in [3] that the category of kG -modules is equivalent to the category of $kQ/\langle Q_{p^a} \rangle$ -modules. In other words the k -algebras kG and $kQ/\langle Q_{p^a} \rangle$ are Morita equivalent. Note that in general they are not isomorphic. A necessary and sufficient condition for the existence of an algebra isomorphism between kG and $kQ/\langle Q_{p^a} \rangle$ is the commutativity of the group K . Since the Hochschild cohomology $H^*(\Lambda, \Lambda)$ is for every k -algebra Λ Morita invariant as a Gerstenhaber algebra (see [6])

p.143), we conclude that the Lie algebras $H^1(kG, kG)$ and $H^1(kQ/\langle Q_{p^a} \rangle, kQ/\langle Q_{p^a} \rangle)$ are isomorphic. Furthermore $H^1(kQ/\langle Q_{p^a} \rangle, kQ/\langle Q_{p^a} \rangle)$ is a product of Lie algebras $H^1(kC/\langle C_{p^a} \rangle, kC/\langle C_{p^a} \rangle)$ where C is a crown of Q , because the quiver Q is a disjoint union of crowns and the Hochschild cohomology is additive on a product of algebras. Thus we have

$$H^1(kG, kG) = \prod_{C \text{ crown of } Q} H^1(kC/\langle C_{p^a} \rangle, kC/\langle C_{p^a} \rangle)$$

Theorem 5.1 *Let k be an algebraically closed field and G a finite group such that the characteristic $p > 0$ of k divides the order of G . Let $n = p^a q$ be the order of G with q prime to p . Suppose that G contains only one Sylow p -subgroup which in addition is cyclic. The following conditions are equivalent:*

- (i) *The Lie algebra $H^1(kG, kG)$ is semisimple.*
- (ii) *The group G is the direct product of a group K of order q and of a cyclic group C_p of order p . The characteristic p of the field k is different from 2.*
- (iii) *$H^1(kG, kG)$ is a product of Witt algebras $W(1, 1) := \text{Der}(k[X]/(X^p))$ and the characteristic p of the field k is different from 2.*

Proof: Let Q be the above-described quiver associated to the group G .

(i) $\Rightarrow$ (ii) : It is clear that the Lie algebra $H^1(kG, kG)$ is semisimple if and only if the Lie algebra $H^1(kC/\langle C_{p^a} \rangle, kC/\langle C_{p^a} \rangle)$ is semisimple for every crown C of the quiver Q . From Proposition 4.2 and Theorem 4.15 we deduce that Q does not have a crown of length ≥ 2 , i.e. Q is a disjoint union of loops. Therefore we have $\chi(x) = 1$ for all $x \in G$ which implies that G is the direct product of the Sylow p -subgroup S and its supplement K . Proposition 4.3 shows that for every loop C of the quiver Q the Lie algebra $H(kC/\langle C_{p^a} \rangle, kC/\langle C_{p^a} \rangle)$ is semisimple if and only if $p^a = p > 2$.

(ii) $\Rightarrow$ (iii) : Let $G = K \times C_p$ and so $\chi = 1$. Therefore Q is a disjoint union of loops. According to Proposition 4.3 we have for every loop C of the quiver Q that $H^1(kC/\langle C_{p^a} \rangle, kC/\langle C_{p^a} \rangle)$ is isomorphic to the Witt Lie algebra $W(1, 1)$.

(iii) $\Rightarrow$ (i) : This is clear, because the Witt algebra is one of the nonclassical simple Lie algebras if the characteristic $p > 0$ of k is different from 2.

q.e.d. 5.1

Corollary 5.2 *Let k be an algebraically closed field and G a finite group such that the characteristic $p > 0$ of k divides the order of G . Let $n = p^a q$ be the order of G with q prime to p . Suppose that G contains only one Sylow p -subgroup which in addition is cyclic. The following conditions are equivalent:*

- (i) *The Lie algebra $H^1(kG, kG)$ is simple.*
- (ii) *The group G is cyclic of order p and the characteristic p of the field k is different from 2.*
- (iii) *$H^1(kG, kG)$ is isomorphic to the Witt Lie algebra $W(1, 1) := \text{Der}(k[X]/(X^p))$ and the characteristic p of the field k is different from 2.*

Proof: Let Q be the above-described quiver associated to the group G .

(i) $\Rightarrow$ (ii) : Let the Lie algebra $H^1(kG, kG)$ be simple and thus semisimple. The preceding Theorem shows that G is the direct product of a group K of order q and of a cyclic group C_p of order p , where q and p are prime. Therefore we have $\chi = 1$ which implies that the quiver Q is a disjoint union of loops. From the fact

that $H^1(kG, kG) = \prod H^1(kC/\langle C_p \rangle, kC/\langle C_p \rangle)$ is a simple Lie algebra it follows that Q has only one loop. Since the field k is algebraically closed and since p does not divide the order q of the group K , the number of isomorphism classes of simple kK -modules, which is the number of vertices of Q , is equal to the number of conjugation classes of the group K . Thanks to the fact that the loop Q has only one vertex we obtain that K has only one conjugation class and so $K = \{1\}$.

(ii) $\Rightarrow$ (iii) : Since the quiver Q associated to the cyclic group C_p is the loop, Proposition 4.3 shows that $H^1(kG, kG)$ is isomorphic to the Witt Lie algebra.

(iii) $\Rightarrow$ (i) : clear

q.e.d. 5.2

References

- [1] M.J. Bardzell, The alternating syzygy behavior of monomial algebras, *J. Algebra* **188**, No. 1 (1997), 69-89.
- [2] C. Cibils, Hochschild cohomology algebra of radical square zero algebras, Algebras and modules II: Eighth International Conference on Representations of Algebras, II (Geiranger, 1996), 93-101, *CMS Conf.Proc.* **24**, Amer. Math. Soc. Providence, RI., 1998.
- [3] P. Gabriel, Problèmes actuels de théorie des représentations, *Enseign. Math.* (2) **20** (1974), 323-332.
- [4] P. Gabriel, A.V. Roiter, Representations of finite-dimensional algebras, with a chapter by B. Keller, *Encyclopaedia of Math. Sci.* **73**, Algebra VIII, Springer, Berlin, 1992.
- [5] M. Gerstenhaber, The cohomology structure of an associative ring, *Ann. of Math.* **78**, No.2 (1963), 267-288.
- [6] M. Gerstenhaber, S.D. Schack, Algebraic cohomology and deformation theory, in "Deformation theory of algebras and structures and applications" (M. Hazewinkel and M. Gerstenhaber, Eds.), NATO Advanced Science Institutes Series C: Mathematical and Physical Sciences, Vol. 247, pp. 11-264, Kluwer Academic Publishers, Dordrecht/Boston, 1988.
- [7] M. Gerstenhaber, S.D. Schack, Relative Hochschild cohomology, rigid algebras, and the Bockstein, *J. Pure Appl. Algebra* **43**, No. 1 (1986), 53-74.
- [8] F. Guil-Asensio, M. Saorín, The group of outer automorphisms and the Picard group of an algebra, *Algebr. Represent. Theory* **2**, No. 4 (1999), 313-330.
- [9] F. Guil-Asensio, M. Saorín, The automorphism group and the Picard group of a monomial algebra, *Comm. Algebra* **27**, No. 2 (1999), 857-887.
- [10] G. Hochschild, On the cohomology groups of an associative algebra, *Ann. of Math.* **46**, No. 2 (1946), 58-67.
- [11] B. Huisgen-Zimmermann, M. Saorín, Geometry of chain complexes and outer automorphisms under derived equivalence, *Trans. Amer. Math. Soc.* **353**, No. 12 (2001), 4757-4777.
- [12] J.E. Humphreys, "Linear algebraic groups", *Graduate Texts in Math.* Vol. 21, Springer, Berlin, 1975.

- [13] B. Keller, Hochschild cohomology and derived Picard groups, Lecture at the international workshop “Representation Theory around the Channel” in Amiens, France, 31 August to 1 September 2001.
- [14] A.C. Locateli, Hochschild cohomology of truncated quiver algebras, *Comm. Algebra* **27**, No. 2 (1999), 645-664.
- [15] R.D. Pollack, Algebras and their automorphism groups, *Comm. Algebra* **17**, No. 8 (1989), 1843-1866.
- [16] W.R. Scott, “Group theory”, Prentice-Hall, Inc., Englewood Cliffs, New Jersey, 1964.
- [17] C.A. Weibel, “An introduction to homological algebra”, *Cambridge Studies in Advanced Mathematics* Vol. 38, Cambridge University Press, Cambridge, 1994.